

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 August 2001 (09.08.2001)

PCT

(10) International Publication Number
WO 01/57202 A2

(51) International Patent Classification⁷: **C12N 15/00**,
C07K 1/00

(21) International Application Number: PCT/US01/03518

(22) International Filing Date: 2 February 2001 (02.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/180,374 4 February 2000 (04.02.2000) US

(71) Applicant: **THE BOARD OF TRUSTEES OF THE
UNIVERSITY OF ARKANSAS** [US/US]; 2404 North
University Avenue, Little Rock, AR 72207-3608 (US).

(72) Inventor: **SHAUGHNESSY, John, D.**; 4317 Old Oak Dr.,
Little Rock, AR 72202 (US).

(74) Agent: **ADLER, Benjamin, A.**; Adler & Associates, 8011
Candle Lane, Houston, TX 77071 (US).

(81) Designated States (*national*): AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with declaration under Article 17(2)(a); without abstract;
title not checked by the International Searching Authority

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/57202 A2

(54) Title: *EV127* GENE SEQUENCES AND PROTEIN ENCODED THEREBY

(57) Abstract:

5 **EVI27 GENE SEQUENCES AND PROTEIN ENCODED THEREBY**

BACKGROUND OF THE INVENTION

10

Cross-Reference to Related Application

 This non-provisional patent application claims benefit of provisional patent application U.S. Serial Number 60/180,374, filed February 4, 2000, now abandoned.

15

Federal Funding Legend

 This invention was produced in part using funds obtained through a grant from the National Cancer Institute. Consequently, the federal government has certain rights in this
20 invention.

Field of the Invention

 The present invention relates generally to the field of molecular biology. More specifically, the present invention
25 relates to the cloning and characterization of a murine and human gene that encodes a novel protein with homology to the IL-17 receptor.

Description of the Related Art

Retroviral insertional mutagenesis in BXH2 and AKXD recombinant inbred (RI) mice induces a high incidence of myeloid leukemia and the proviral integration sites in the leukemias provide powerful genetic tags for disease gene identification (Bedigian et al., 1984; Gilbert et al., 1993). During the past several years, a number of disease genes have been identified in these leukemias by proviral tagging. These disease genes include a tumor suppressor gene, *neurofibromatosis type 1* (*Nf1*); a gene with homology to the lymphoid-restricted type II membrane protein *Jaw1*, *Mrv integration site 1* (*Mrvil*); a gene encoding a hematopoietic cell growth and differentiation factor, *myeloblastosis oncogene* (*Myb*); three homeobox genes, *homeobox A7* (*Hoxa7*), *homeobox A9* (*Hoxa9*), and *myeloid ecotropic viral integration site 1* (*Meis1*); a zinc-finger protein (*Evi1*); and a gene with homology to the *ubiquitin-specific protease 8* (*Usp8*) oncogene and to genes encoding various cell cycle regulatory proteins, *ecotropic viral integration site 5* (*Evi5*) (Buchberg et al., 1990, Viskochil et al., 1990, Shaughnessy et al., 1999; Copeland and Jenkins, 1999, Nakamura et al., 1996a, Morishita et al., 1988, Liao, et al., 1997). Four of the genes are proven or suspected human disease genes: *EVII*, *NF1* and *HOXA9* are causally associated with myeloid leukemia and *EVI5* with stage 4S neuroblastoma (Ogawa et al., 1996, Copeland and Jenkins, 1999, Nakamura et al., 1996b, Roberts et al. 1998), validating the usefulness of this approach for human disease gene identification.

Although proviral tagging has identified many disease genes, it is apparent that several more genes remain to be cloned. This is suggested by the fact only 45% of BXH2 leukemias contain a virally induced mutation in one of the genes identified so far.

5 Disease genes for 55% of BXH2 leukemias remain to be identified. The same is true for human acute myeloid leukemias (AMLs) where the 11 different chromosomal translocations and inversions cloned to date are found in only 45% of acute myeloid leukemias (Look, 1997). Disease genes for 55% of acute myeloid

10 leukemias remain to be identified. Ultimately, it should be possible to use proviral tagging to do a saturation screen for BXH2 disease genes. The expectation is that some of these genes will represent human acute myeloid leukemias genes that are not easy to clone because they are infrequently involved in human

15 disease or are not marked by a cytologically detectable rearrangement. Given the large number of genes that may remain to be identified, this task could be difficult using conventional proviral tagging approaches, which rely on cloning leukemia-specific proviral integration sites into bacteriophage lambda.

20 With this potential problem in mind, an inverse PCR (IPCR) method for proviral tagging was developed that makes use of automated DNA sequencing and the genetic tools provided by the Mouse Genome Project, which greatly increases the throughput of proviral tagging for disease gene identification.

25 More than 400 proviral integration sites from BXH2 myeloid leukemias (and AKXD T- and B-cell leukemias) were cloned and characterized using this inverse PCR method (Li et al., 1999), which lead to the identification of more than 90 new candidate

leukemia disease genes (Li et al., 1999). Nineteen new common integration sites (sites that are targets of viral integration in more than one leukemia) were also identified in these studies and BLAST search and/or chromosome mapping identified candidate
5 disease genes for 12 of these common sites (Li et al., 1999).

One common integration site identified by the inverse PCR is *Evi27* (Li et al., 1999). While BLAST searches did not identify a candidate disease gene for *Evi27*, chromosome mapping studies showed that *Evi27* maps to mouse chromosome 14 in a
10 region of human 3p21 homology (Copeland et al., 1993). This result is interesting because treatment-related 3p21 breaks are often observed in myelodysplastic syndrome (MDS) and AML patients (Shi et al., 1996) and 3p21 is the most frequently deleted region seen in chronic myeloid leukemia (CML)
15 (Johansson et al., 1997). *Evi27* may be, therefore, an important human disease gene.

The prior art is deficient in lack of the characterization a novel cytokine receptor-related gene whose expression is upregulated by viral integration at *Evi27*. The
20 present invention fulfills this long-standing need in the art.

SUMMARY OF THE INVENTION

The present invention describes the cloning and
25 sequences of a novel IL-17 receptor-related gene in human and mouse whose expression is upregulated by viral integration in a murine acute myeloid leukemia. Gene transcription and protien expression were examined by northern blot analysis, western blot

analysis and immunohistochemical staining. The gene disclosed herein may facilitate myeloid cell transformation and be involved in human disease.

5 **BRIEF DESCRIPTION OF THE DRAWINGS**

The appended drawings have been included herein so that the above-recited features, advantages and objects of the invention will become clear and can be understood in detail.

10 These drawings form a part of the specification. It is to be noted, however, that the appended drawings illustrate preferred embodiments of the invention and should not be considered to limit the scope of the invention.

Figure 1 shows a partial long-range restriction map
15 of the *Evi27* locus. Bacteriophage λ and P1 clones used for gene localization and exon trapping are shown above the map. The location of the 5' end and transcriptional orientation of the *Evi27* and *Cdhd* genes are noted by arrows above the map. The location of the exon with homology to the yeast *ARP8* gene is also noted.
20 The transcriptional orientation of this gene has not been determined. The position and transcriptional orientation of the proviruses in the B160 and 15-38374 leukemias are indicated below the map. The cluster of rare cutting restriction enzymes surrounding the proviral integration sites is also noted. The 14kb
25 *Bam*HI fragment cloned from leukemia B160 is presented below the map. The viral sequences and viral long terminal repeat (LTR) is noted by a thick line and black box, respectively. The position of the 5' end of *Evi27* with respect the viral LTR is noted.

Restriction enzymes: B, *Bam*HI; Bs, *Bss*HII; C, *Cla*I; E, *Eag*I; M, *Mlu*I, N, *Nae*I; Nr, *Nru*I, S, *Sac*II; Sa, *Sal*I;

Figure 2 shows Northern blot analyses of *Evi27* expression in murine tissues and cancer cell lines. Multiple tissue Northern blots of normal adult (**Figure 2A**), BXH2 leukemic cell lines (**Figure 2B**), and various hematopoietic cell lines (**Figure 2C**): M1, myeloid leukemia; WEHI3B, monocyte; HYB. 548, B-cell hybridoma; WEHI231, preB-cell lymphoma; R1.1, lymphocytic thymoma; EL4, thymoma, P815, mastocytoma. The blots were also hybridized with a GAPDH or β -actin probe to control for RNA loading. The size in kilobases (kb) of molecular weight markers is show on the left of each panel. To the right of each panel are the sizes of the *Evi27* transcripts observed

Figure 3 shows Northern blot analyses of *EVI27* expression in human tissues and cancer cell lines. Multiple tissue Northern blots of normal adult (**Figure 3A**), immune tissues (**Figure 3B**) and cancer cell lines (**Figure 3C**) hybridized with an EST specific for the human *EVI27* gene. The human cancer cell lines are HL60, promyelocytic leukemia; HeLa, cervical carcinoma; K562, chronic myelogenous leukemia; MOLT4, T-lymphoblastic leukemia; Raji, Burkitt's lymphoma; SW480, colon adenocarcinoma; A549, lung carcinoma; G361, melanoma. The blots were also hybridized with a β -actin probe to control for RNA loading. The size in kb of molecular weight markers is show on the left of each panel.

Figure 4 shows an alignment of mouse and human *Evi27* proteins. The putative signal peptide, transmembrane domain, and peptide used to generate polyclonal antisera are

indicated by a black line over the sequence. Conserved protein motifs are indicated by a double line. Conserved amino acids are boxed with amino acid identities noted in bold. Normal typeface letters represents conservative amino acid changes. Gaps created
5 in the sequences to optimize alignments are indicated by a dash in the sequence string. Abbreviations: N-GLY, N-linked glycosylation; GSK, kinase phosphorylation site.

Figure 5 shows the Evi27 gene structure. The gene consists of 11 exons with the first and last exons containing
10 untranslated regions. The transmembrane domain is located in exon 10.

Figure 6 shows an alignment of the human EVI27 and IL-17R proteins. Conserved amino acids are boxed with amino acid identities noted in bold. Gaps created in the sequences to
15 optimize alignments are represented by dashes. Amino acid positions are indicated to the right and left of the sequence.

Figure 7 shows the chromosome mapping of the human *EVI27* gene. **Figure 7A** fluorescence *in situ* hybridization of a PAC clone specific for *EVI27* gene to normal human
20 metaphase chromosomes. Arrows indicate localization of the probe. **Figure 7B** shows G-banding of the same metaphase spread in (**Figure 7A**) following destaining of FISH reagents. The probe is localized to band 3p21.

Figure 8 shows immunofluorescence and Northern
25 blot analysis of *Evi27* expression in murine cell line. The 2.7 kb cDNA coding for the 55 kD isoform of the membrane bound form of the murine Evi27 gene was cloned into a eukaryotic expression plasmid vector and transfected into the murine myeloid leukemia

cell line 32D. A cell line was established by limiting dilution and called 32DEvi27A. **Figure 8A** shows a Northern blot hybridization of poly-A mRNA from 32D and 32D/Evi27A stable transfectants. Note the abundant expression of the transgene in the transfectant and the absence of expression in the parental line. The blot was also hybridized with a β -actin probe to control for RNA loading. The size in kb of molecular weight markers is shown on the left of each panel. **Figure 8B** shows Western blot analysis of same cell lines with affinity purified anti-Evi27 antisera. The transfectant shows overexpression of a 55 kD protein as expected from predicted mRNA translation of the 2.7 kb Evi27 cDNA. Markers in kD are to the left. **Figure 8C** shows immunofluorescence staining of the Evi27 protein (red) in the myeloid cell line 32D (top panel) and 32D cells transfected with an Evi27 cDNA expression construct (lower panel). Note the light staining in the parental line and abundant staining in the transfectant (left panels). Nuclei are stained blue with DAPI. Cells were also stained with Evi27 antibody preincubated with Evi27 peptide (right panels). Note that no red staining is evident, demonstrating specificity of the antisera. **Figure 8D** shows cell surface expression of Evi27 by flow activated cell sorting analysis (red: anti-Evi27; blue: anti-Evi27+peptide; green: IgG control).

Figure 9 shows cell surface expression and capping of Evi27 on murine T cell lymphoma EL-4. The left panel shows immunochemistry of the EL-4 cells with the anti-Evi27 antibody before and after pre-treatment with Evi27 peptide. Right panel shows FACS analysis with increasing amount of anti-Evi27 antibody. The capping of Evi27 possibly indicates

polydimerization and receptor activation of Evi27.

Figure 10 shows subcellular localization of Evi27 in myeloid leukemia B187 and B190. The left panel shows immunochemistry of the B187 and B190 cells with the anti-Evi27 antibody before and after pre-treatment with Evi27 peptide. B187 cells show diffuse speckled staining in the cytoplasm, whereas B190 cells shows a capping pattern similar to the EL-4 cells. The signals could be competed away by preincubation with Evi27 peptide. Right panel shows western blot analysis of same cell lines with anti-Evi27. Both cell lines express multiple isoforms of Evi27, and B187 cells express more of the 55 kD isoform. The bands could be competed away by preincubation with Evi27 peptide.

Figure 11 shows abundant Evi27 protein was found in the cytoplasm, but not on the cell surface, in the myeloid leukemia B160. The left panel shows immunochemistry of the B160 cells with the anti-Evi27 antibody before and after pre-treatment with Evi27 peptide. Evi27 is expressed in the cytoplasm at high concentration. Right panel shows FACs analysis with anti-Evi27, indicating Evi27 is not expressed on the cell surface (red: anti-Evi27; black: IgG control).

Figure 12 shows Western blot analysis of Evi27 protein expression in BXH2 leukemia cell lines, EL4 and R1.1 T-cell lines, and the WEHI231 B-cell line. 10 µg of protein from cell lysates was run in tandem and hybridized with Evi27 antibody either without (**Figure 12A**) or with (**Figure 12B**) preincubation with Evi27 peptide. Note the absence or reduction of specific bands in panel (**Figure 12B**) compared to (**Figure 12A**).

Molecular weight standards in kD are indicated to the left. The Evi27 isoforms and sizes are indicated to the right of panel A.

Figure 13 shows western blot analysis of Evi27 expression in human cell lines. SW480 and SW620 are colon adenocarcinoma; MEG01, megakaryocytic leukemia; THP-1, monocytic leukemia; K562, chronic myelogenous leukemia; HL-60, promyelocytic leukemia; U266, ARP1 and 8226, multiple myeloma; HeLa, cervical carcinoma; HEL, erythrocytic leukemia. 55 kD and 30 kD are major bands recognized, thus the antibody recognizes both human Evi27 protein isoforms. The bands could be competed away by preincubation of Evi27 antisera with Evi27 peptide as shown in the right panel. MEG01, K562 and HEL cell lines express the highest levels of the Evi27 protein.

Figure 14 shows surface expression of Evi27 on human hematopoietic cell lines. Left panel shows FACs analysis of Evi27 expression on the surface of various human cell lines (red: affinity purified anti-Evi27 antisera; black: IgG control). Evi27 was expressed on THP-1 and K562 cells but not the HL-60 cells. Right panel shows western blot analysis of same cell lines with anti-Evi27. As expected from the FACs data, THP-1 cells express more Evi27 protein than K562 cells, which in turn express more Evi27 than HL-60 cells. It is also shown that K562 cells produce more of the 30 kD soluble form of the Evi27 protein.

DETAILED DESCRIPTION OF THE INVENTION

Evi27 is a common site of retroviral integration in BXH2 murine myeloid leukemias. The present invention shows

that integration at *Evi27* occurs in a CpG island ~6 kb upstream from a novel gene (designated *Evi27*) with homology to the *IL17 receptor (Il17r)* and that proviral integrations result in increased *Evi27* expression. The human *EVI27* homologue was also cloned and mapped to chromosome 3p21. Multiple *Evi27* isoforms were detected at the RNA and protein level in both human and mouse, indicating that *Evi27* expression is complex. Some of the isoforms are shown to likely represent secreted soluble forms of the protein produced by intron incorporation or by proteolytic cleavage. In the mouse, highest *Evi27* expression occurs in liver and testes with lower expression in kidney and lung. In humans, *EVI27* is expressed at high levels in the kidney, with moderate levels in the liver, brain, and pancreas. Within hematopoietic cells, *Evi27* expression is restricted. Northern and Western analysis showed that *Evi27* is expressed in selected T-cell, B-cell and myeloid cell lines. These results suggest that *Evi27* expression is tightly regulated during hematopoietic differentiation. Collectively, these studies identify a new member of the cytokine receptor family whose increased and uncoordinated expression may lead to myeloid leukemia by altering *Evi27's* normal ability to control the growth and/or differentiation of hematopoietic cells.

In accordance with the present invention, there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook, Fritsch & Maniatis, "Molecular Cloning: A Laboratory Manual (2nd. Ed.)", (1989); "DNA Cloning: A Practical Approach,"

Volumes I and II (D.N. Glover ed. 1985); "Oligonucleotide Synthesis" (M.J. Gait ed. 1984); "Nucleic Acid Hybridization" [B.D. Hames & S.J. Higgins eds. (1985)]; "Transcription and Translation" [B.D. Hames & S.J. Higgins eds. (1984)]; "Animal Cell Culture" [R.I. Freshney, ed. (1986)]; "Immobilized Cells And Enzymes" [IRL Press, (1986)]; B. Perbal, "A Practical Guide To Molecular Cloning" (1984). Therefore, if appearing herein, the following terms shall have the definitions set out below.

A "DNA molecule" refers to the polymeric form of deoxyribonucleotides (adenine, guanine, thymine, or cytosine) in its either single stranded form or a double-stranded helix. This term refers only to the primary and secondary structure of the molecule, and does not limit it to any particular tertiary forms. Thus, this term includes double-stranded DNA found, *inter alia*, in linear DNA molecules (*e.g.*, restriction fragments), viruses, plasmids, and chromosomes. DNA structures are discussed herein according to the normal convention of giving only the sequence in the 5' to 3' direction along the nontranscribed strand of DNA (*i.e.*, the strand having a sequence homologous to the mRNA). The term "peptide" or "polypeptide" is defined as several (*i.e.*, multiple) amino acids attached together.

A "vector" is a replicon, such as plasmid, phage or cosmid, to which another DNA segment may be attached so as to bring about the replication of the attached segment. A "replicon" is any genetic element (*e.g.*, plasmid, chromosome, virus) that functions as an autonomous unit of DNA replication *in vivo*; *i.e.*, capable of replication under its own control. An "origin of replication" refers to those DNA sequences that participate in or

direct DNA synthesis. An "expression control sequence" is a DNA sequence that controls and regulates the transcription and translation of another DNA sequence. A coding sequence is "operably linked" and "under the control" of transcriptional and translational control sequences in a cell when RNA polymerase transcribes the coding sequence into mRNA, which is then translated into the protein encoded by the coding sequence.

In general, expression vectors containing promoter sequences which facilitate the efficient transcription and translation of the inserted DNA fragment are used in connection with a particular host. The expression vector typically contains an origin of replication, promoter(s), terminator(s), as well as specific genes which are capable of providing phenotypic selection in transformed cells (*i.e.*, selectable markers). The transformed hosts can be fermented and cultured according to means known in the art to achieve optimal cell growth.

A DNA "coding sequence" is a double-stranded DNA sequence which is transcribed and translated into a polypeptide *in vivo* when placed under the control of appropriate regulatory sequences. Specifically as used herein, "DNA coding for a protein" means DNA sequences which produce a particular primary amino acid sequence. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxyl) terminus. A coding sequence can include, but is not limited to, prokaryotic sequences, a cDNA from eukaryotic mRNA, genomic DNA sequences from eukaryotic (*e.g.*, mammalian) organisms and even synthetic DNA sequences. A polyadenylation signal and

transcription termination sequence will usually be located 3' of the coding sequence. A "cDNA" is defined as copy DNA or complementary DNA and is a product of a reverse transcription reaction from an mRNA transcript. An "exon" is an expressed
5 sequence transcribed from the gene locus, whereas an "intron" is a non-expressed sequence that is from the gene locus.

Transcriptional and translational control sequences are DNA regulatory sequences, such as promoters, enhancers, polyadenylation signals, terminators, and the like, that provide
10 for the expression of a coding sequence in a host cell. A "cis-element" is a nucleotide sequence, also termed a "consensus sequence" or "motif", that interacts with proteins that upregulate or downregulate expression of a specific gene locus. A "signal sequence" can also be included with the coding sequence. This
15 sequence encodes a signal peptide, N-terminal to the polypeptide, that communicates to the host cell and directs the polypeptide to the appropriate cellular location. Signal sequences can be found associated with a variety of proteins native to prokaryotes and eukaryotes.

20 A "promoter sequence" is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. Specifically as used herein, the term "promoter(s)" means regulatory DNA sequences that control transcription of the cDNA.
25 For purposes of defining the present invention, a minimal promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary

to initiate transcription at levels detectable above background. Within the promoter sequence will be found a transcription initiation site, as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase. 5 Eukaryotic promoters often, but not always, contain "TATA" boxes and "CAT" boxes. Prokaryotic promoters contain the -10 and -35 consensus sequences, and additionally, ribosomal binding Shine-Dalgarno sequences. As used herein, "promoter" may also refer to an intact regulatory sequence directing transcription 10 (and subsequent translation) of a coding sequence, and may include any or all of the above-mentioned transcriptional and translational control sequences.

The term "oligonucleotide" is defined as a molecule comprised of two or more deoxyribonucleotides, preferably more 15 than three. Its exact size will depend upon many factors which, in turn, depend upon the ultimate function and use of the oligonucleotide. The term "primer" as used herein refers to an oligonucleotide, whether occurring naturally as in a purified restriction digest or produced synthetically, which is capable of 20 acting as a point of initiation of synthesis when placed under conditions in which synthesis of a primer extension product, which is complementary to a nucleic acid strand, is induced, *i.e.*, in the presence of nucleotides and a polymerizing agent, such as a DNA polymerase, and at a suitable temperature and pH. The 25 primer may be either single-stranded or double-stranded and must be sufficiently long to prime the synthesis of the desired extension product in the presence of the polymerizing agent. The exact length of the primer will depend upon many factors,

including temperature, source of primer and use the method. For example, for diagnostic applications, depending on the complexity of the target sequence, the oligonucleotide primer typically contains 15-25 or more nucleotides, although it may
5 contain fewer nucleotides.

“Recombinant DNA technology” refers to techniques for uniting two heterologous DNA molecules, usually as a result of *in vitro* ligation of DNAs from different organisms. Recombinant DNA molecules are commonly produced by
10 experiments in genetic engineering. Synonymous terms include “gene splicing”, “molecular cloning” and “genetic engineering”. The product of these manipulations results in a “recombinant” or “recombinant molecule”.

A cell has been “transformed” or “transfected” with
15 exogenous or heterologous DNA when such DNA has been introduced into the cell. Specifically as used herein, the term “transformation” or “transfection” means incorporation permitting expression of heterologous DNA sequences by a cell. The transforming DNA may or may not be integrated (covalently
20 linked) into the genome of the cell. In prokaryotes, yeast, and mammalian cells, the transforming DNA may be maintained on an episomal element such as a vector or plasmid. With respect to eukaryotic cells, a stably transformed cell is one in which the transforming DNA has become integrated into a chromosome so
25 that it is inherited by daughter cells through chromosome replication. This stability is demonstrated by the ability of the eukaryotic cell to establish cell lines or clones comprised of a population of daughter cells containing the transforming DNA. A

"clone" is a population of cells derived from a single cell or ancestor by mitosis. A "cell line" is a clone of a primary cell that is capable of stable growth *in vitro* for many generations. An organism, such as a plant or animal, that has been transformed
5 with exogenous DNA is termed "transgenic".

As used herein, the term "host" is meant to include not only prokaryotes, but also eukaryotes such as yeast, plant and animal cells. Specifically as used herein, the term "host(s)" means any cell that will allow or direct expression. Specifically as
10 used herein, "chimeric cell" means a cell whose DNA has been altered compared to a normal cell of the same organism. A recombinant DNA molecule or gene can be used to transform a host using any of the techniques commonly known to those of ordinary skill in the art. Prokaryotic hosts may include *E. coli*, *S.*
15 *tymphimurium*, *Serratia marcescens* and *Bacillus subtilis*. Eukaryotic hosts include yeasts such as *Pichia pastoris*, mammalian cells and insect cells, and plant cells, such as *Arabidopsis thaliana* and *Tobaccum nicotiana*.

As used herein, "fragment," as applied to a
20 polypeptide, will ordinarily be at least 10 amino acid residues, more typically at least 20 residues, and preferably at least 30 (*e.g.*, 50) residues in length, but less than the entire, intact sequence. Fragments can be generated by methods known to those skilled in the art, *e.g.*, by enzymatic digestion of naturally
25 occurring or recombinant protein, by recombinant DNA techniques using an expression vector that encodes a defined fragment, or by chemical synthesis. Purified fragments or antigenic fragments can be used to generate antibodies employing

standard protocols known to those skilled in the art. As used herein, "functional fragment" is meant to encompass those peptide fragments retaining biological activity of Evi27.

Due to the redundancy of the DNA code, there are
5 millions of DNA sequences that would produce the same amino acid sequence when expressed. Given an amino acid sequence, one can substitute into the natural DNA sequence alternative codons for the desired amino acids to produce an alternative DNA sequence also coding for the novel protein. One may find
10 that particular chimeric cells of a particular expression method favor particular mRNA codons for a particular amino acid. Altering the human DNA sequence to increase the frequency of favored codons may improve the expression efficacy in a chimeric cell, thus improving the efficacy of the expression
15 process. The sequences may be derived by substitution of redundant codons for the amino acid sequences and splicing the substituted sequences into the natural gene by routine methods well known in the art. Those skilled in the art will recognize that many variations are possible in substituting conserved amino
20 acids in the protein sequence which will produce variations in sequence without seriously changing the biological activity of the protein. It is impractical to attempt to list all the millions of DNA sequences that may code for the claimed sequence. However, the invention comprises the novel protein, its novel amino acid
25 sequence, and all DNA sequences natural or synthetic coding for the novel amino acid sequence.

These substitution analogs may be constructed in the following manner: Table 1 lists the alternative codons that code

for the 20 common amino acids. DNA sequence substitution analogs that also code for human can be constructed by choosing alternate codons from Table 1 to alter the DNA sequence between a pair of restriction enzyme cleavage sites, as are well known in the art. Alternative codons are assembled into a synthetic oligonucleotide by conventional methods and the synthetic oligo is substituted into the endonuclease treated DNA by the methods described in "Molecular Cloning. A Laboratory Manual", 2d Edition, Cold Spring Harbor Laboratory Press (1989), to produce a substitution analog. Other methods generally known to those skilled in the art can also be employed to obtain substitution analogs of DNA sequences.

The alteration of the DNA by cleavage and codon substitution may be repeated to substitute substantial portions of the original DNA sequence with alternative codons without altering the protein amino acid sequence. Alteration of a DNA sequence which produces no change in the protein expressed by the DNA sequence might, for example, be conducted to increase protein expression in a particular host cell by increasing the occurrence of codons that correspond to amino acid tRNAs found in higher concentration in the host cell. Such altered DNA sequences for substitution analogs can be easily produced by those of ordinary skill in the art following the method set out above, or other alternative techniques for altering the DNA sequence while obtaining the same protein on expression. Substitution analogs can be obtained by substitution of oligonucleotides at restriction cleavage sites as described above, or by other equivalent methods that change the codons while

preserving the amino acid sequence of the expressed protein.

TABLE 1

<u>SYMBOL</u>		<u>AMINO ACID</u>	<u>CODON USAGE</u>
5	<u>1-Letter</u> <u>3-Letter</u>		
	A Ala	Alanine	GCT, GCC, GCA, GCG
	C Cys	Cysteine	TGT, TGC
	D Asp	Aspartic acid	GAT, GAC
	E Glu	Glutamic acid	GAA, GAG
10	F Phe	Phenylalanine	TTT, TTC
	G Gly	Glycine	GGT, GGC, GGA, GGG
	H His	Histidine	CAT, CAC
	I Ile	Isoleucine	ATT, ATC, ATA
	K Lys	Lysine	AAA, AAG
15	L Leu	Leucine	TTA, TTG, CTT, CTC, CTA, CTG
	M Met	Methionine	ATG
	N Asn	Asparagine	AAT, AAC
	P Pro	Proline	CCT, CCC, CCA, CCG
	Q Gln	Glutamine	CAA, CAG
20	R Arg	Arginine	CGT, CGC, CGA, CGG, AGA, AGG
	S Ser	Serine	TCT, TCC, TCA, TCG, AGT, AGC
	T Thr	Threonine	ACT, ACC, ACA, ACG
	V Val	Valine	GTT, GTC, GTG, GTG
	W Trp	Tryptophan	TTG
25	Y Tyr	Tyrosine	TAT, TAC

As described herein, a standard Northern blot assay can be used to ascertain the relative amounts of mRNA in a cell or tissue in accordance with conventional Northern hybridization techniques known to those persons of ordinary skill in the art.

5 Alternatively, a standard Southern blot assay may be used to confirm the presence, the copy number and/or the position of a gene in accordance with conventional Southern hybridization techniques known to those of ordinary skill in the art. Both the Northern blot and Southern blot use a hybridization probe, (*e.g.*,

10 radiolabelled full-length or partial cDNA) of at least 20 (preferably at least 30, more preferably at least 50, and most preferably at least 100) consecutive nucleotides in length. The DNA hybridization probe can be labelled by any of the many different methods known to those skilled in this art.

15 The labels most commonly employed for these studies are radioactive elements, enzymes, chemicals which fluoresce when exposed to ultraviolet light, and others. A number of fluorescent materials are known and can be utilized as labels. These include, for example, fluorescein, rhodamine, auramine,

20 Texas Red, AMCA Blue and Lucifer Yellow. Proteins can also be labeled with a radioactive element or with an enzyme. The radioactive label can be detected by any of the currently available counting procedures. The preferred isotope may be selected from ^3H , ^{14}C , ^{32}P , ^{35}S , ^{36}Cl , ^{51}Cr , ^{57}Co , ^{58}Co , ^{59}Fe , ^{90}Y , ^{125}I , ^{131}I , and

25 ^{186}Re .

Enzyme labels are likewise useful, and can be detected by any of the presently utilized colorimetric, spectrophotometric, fluorospectrophotometric, amperometric or gasometric

techniques. The enzyme is conjugated to the selected particle by reaction with bridging molecules such as carbodiimides, diisocyanates, glutaraldehyde and the like. Many enzymes which can be used in these procedures are known and can be utilized.

5 The preferred are peroxidase, β -glucuronidase, β -D-glucosidase, β -D-galactosidase, urease, glucose oxidase plus peroxidase and alkaline phosphatase. U.S. Patent Nos. 3,654,090, 3,850,752, and 4,016,043 are referred to by way of example for their disclosure of alternate labeling material and methods.

10 The present invention is directed to an isolated nucleic acid molecule encoding an IL-17 receptor-related protein selected from the group consisting of: (a) an isolated nucleic acid molecule of SEQ. ID NO: 1, 2, 3 or 4 which encodes an IL-17 receptor-related protein; (b) an isolated nucleic acid molecule
15 which is complimentary and hybridizes to the nucleic acid molecules of (a); and (c) isolated nucleic acid molecule differing from the isolated nucleic acid molecules of (a) or (b) in codon sequence due to the degeneracy of the genetic code, and which encodes an IL-17 receptor-related protein.

20 In one embodiment, there is provided a fragment of one of the nucleic acid molecules listed above that is at least 10 bases long and which will selectively hybridize to nucleic acid molecule encoding an IL-17 receptor-related protein. Preferably, the nucleic acid molecule may be used as an anti-sense molecule
25 to inhibit the expression of an IL-17 receptor-related protein, for chromosomal mapping or mutation analysis of gene encoding an IL-17 receptor-related protein.

The present invention is also directed to a genomic

DNA encoding an IL-17 receptor-related protein, wherein said genomic DNA hybridizes to the nucleic acid molecules described above.

5 The present invention is also directed to vector comprising a nucleic acid molecule of SEQ. ID NO: 1, 2, 3 or 4 and cells transfected with the vector. Representative cells include bacterial cells, mammalian cells, plant cells and insect cells.

The present invention is also directed to an IL-17 receptor-related protein, or a peptide derived thereof, encoded
10 by nucleic acid molecule of SEQ ID No. 3 or 4, wherein said protein is about 24, 33, 56, 47, 75, 127, and 150 kD in size as detected by western blot analysis in BXH2 leukemia cell. The present invention is further directed to an IL-17 receptor-related protein, or a peptide derived thereof, encoded by the nucleic acid
15 molecules disclosed herein, wherein said protein has the amino acid sequence of SEQ. ID NO: 5, 6, 7 or 8. Preferably, the peptide is at least 4 amino acids long. The peptides can be used to generate anti-Evi27 antibody, and a person having ordinary skill in the art would be readily able to prepare an antibody that binds
20 specifically to any of these proteins or peptides.

The present invention is also directed to a method of stimulating the secretion of cytokines from a cell, comprising the step of binding a ligand to the IL-17 receptor-related protein, Evi27. In general, Evi27 stimulation can lead to the secretion of
25 IL-1, IL-8 and TNF- α from hematopoietic cells, leukemia cells and kidney cells.

The present invention is also directed to a method of modulating the expression and activity of an IL-17 receptor-

related protein, Evi27, comprising the step of contacting a molecule to a cell, wherein the binding of said molecule to Evi27 mRNA or protein results in increased or decreased expression and activity of the Evi27 protein. Preferably, the molecule can be
5 anti-sense oligonucleotides, small molecule that binds to Evi27, modified IL-17E or soluble form of the Evi27 receptor.

The following examples are given for the purpose of illustrating various embodiments of the invention and are not meant to limit the present invention in any fashion:

10

EXAMPLE 1

BXH2 Mice and Leukemic Cell Lines

The BXH2 recombinant inbred strain was obtained
15 from the Jackson Laboratory (Bar Harbor, ME) and maintained at the NCI-Frederick Cancer Research and Development Center. BXH2 leukemia cell lines have been previously described (Largaespada et al., 1995). Other cell lines were purchased from the ATCC and grown in either RPMI 1640 or DMEM supplemented
20 with 2mM glutamine, 4.5 g/L glucose, Pen-Strep, and 10% fetal bovine serum (Atlanta Biologicals, Atlanta GA).

EXAMPLE 2

DNA Extraction and Southern Blot Hybridization

25 High molecular weight genomic DNAs were extracted from frozen normal tissues and leukemic spleens and lymph nodes as previously described (Jenkins et al., 1982). Bacteriophage and plasmid DNAs were purified using standard

procedures (Sambrook et al., 1987). Restriction enzyme digestions, agarose gel electrophoresis, Southern blot transfers, hybridizations, and washes were performed as previously described (Sambrook et al., 1987).

5

EXAMPLE 3

Genomic Cloning

70 μ g of DNA from leukemia N57 was digested to completion with *Bam*HI and fractionated by electrophoresis in TAE buffer. Fragments approximating the somatically acquired pAKV5 (ecotropic murine leukemia virus) hybridizing fragments were cut from the gel and purified using Qiaex beads according to the manufacturer's instructions (Qiagen, Valencia, CA). The *Bam*HI fragments were cloned into EMBL4 phage arms (Stratagene, La Jolla, CA). Ligated material was packaged with Gigapack Gold packaging extracts (Stratagene, La Jolla, CA). Library screening was performed according to the protocol of Benton and Davis (1977). The P1 and lambda genomic clones were derived from commercial libraries (Genome Systems, St. Louis, MO, Clontech, Palo Alto, CA).

EXAMPLE 4

Human Chromosome Mapping

Metaphase chromosomes from normal peripheral blood samples were prepared using conventional methods. Slides were incubated in 2XSSC (pH 7.0, 37°C) for 15 minutes then immersed in 0.1N HCl/0.05% Triton X-100 for 15 minutes at

room temperature (RT). Slides were washed in 2XSSC (pH 7.0) twice at room temperature, then washed in 1XPBS, pH 7.2 once at room temperature. Slides were immersed in 1% formaldehyde (diluted in PBS) for 10 minutes at room temperature, washed
5 twice in 1XPBS at room temperature, then once in 2XSSC (pH 7.0) at room temperature. Chromosomes were dehydrated in an ethanol series and air dried at room temperature.

A human PAC clone specific for *EVI27* was isolated by screening a human P1 artificial chromosome (PAC) library
10 according to the manufacturer's instructions and high molecular weight PAC DNA was isolated using the KB-100 kit (Genome Systems, St. Louis, MO). Spectrum-Red-dUTP (Vysis, Downer's Grove, IL) was incorporated into PAC/BAC probes by nick translation according to the manufacturer's protocol (Vysis,
15 Downer's Grove, IL). DNA was resuspended in Hybrisol VII (Oncor, Gaithersburg, MD) (50% formamide/2xSSC) at a final concentration of 5 ng/ul and hybridized to chromosomes according to manufacturer's instructions. The chromosomes and probe solution was incubated at 75°C for 10 minutes. then to 37°C
20 for 16 hrs in a humidified chamber. Post hybridization washes consisted of 65% formamide /2xSSC (pH 7.0) for 15 minutes at 43°C, then 2XSSC (pH 7.0) for 8 minutes at 37°C, 1XPBD (Oncor, Gaithersburg MD) twice at room temperature, then in 1XPBS (pH 7.2). Finally, 10 µl of propidium iodide/antifade (1:20) was
25 added on each area and a coverslip added. Following FISH, the slide was washed twice in Xylene to remove the coverslip and excess oil. The slide was then washed in 1XPBD at 37°C for 5 minutes followed by a dehydration step through an ethanol

series. The slide was then incubated in 50% formamide/2XSSC for 90 minutes at 37°C, washed briefly in tap water, followed by dehydration in 10, 80, and 95% ethanol and stained with Wright's stain/phosphate buffer for 5 minutes.

5

EXAMPLE 5

Exon Trapping

The P1 clone p57P1 was digested to completion with *Bgl*II and *Bam*HI and purified by phenol extractions and ethanol precipitations. Insert DNA was then ligated to a *Bam*HI digested pSPL3 vector (Life Technologies, Gaithersburg, MD). DNA from the ligations was used to transform DH10B maximum efficiency cells (Life Technologies, Gaithersburg, MD). Additional steps in the experiment were performed exactly according to the manufacturer's protocol (Life Technologies). DNA sequence of approximately 100 individual clones from each ligation was generated as described below.

10
15

EXAMPLE 6

Poly(A)+ RNA Isolation and Northern Blot Analysis

20

Premade Northern blots containing 2 µg of twice selected poly(A)+ RNA from various normal tissues and cell lines were purchased from commercial source (CLONTECH, Palo Alto, CA). Total RNA was extracted from cell line suspensions by the RNAzol B method (Tel-Test, Friendswood, TX). poly(A)+ RNA was purified from the total RNA preps by oligo-d(T) column chromatography according the manufacturers recommendations (Amersham Pharmacia Biotech, Piscataway, NJ). 2-5 µg of

25

poly(A)+ RNA was fractionated by electrophoresis in 1.0 % agarose gels containing formaldehyde and transferred to Hybond N+ membranes (Amersham Pharmacia Biotech, Piscataway, NJ). The membranes were prehybridized and hybridized according to the method of Church and Gilbert or using ExpressHyb solution (Clontech, Palo Alto, CA). Blots were then exposed to X-ray film at -70°C with an intensifying screen.

EXAMPLE 7

cDNA Cloning

cDNA cloning was carried out by a combination of 5' and 3' rapid amplification of cDNA ends (5' and 3' RACE) and modified RT-PCR as described (Shaughnessy et al., 1999). Briefly, the nucleotide sequence of the trapped exon p57P1ET47 was used to design specific nested oligonucleotides. The primary 5' and 3' RACE reactions were performed using mouse liver Marathon Ready cDNA (Clontech, Palo Alto, CA) as a template according to the manufactures protocol using the Advantage cDNA amplification kit (Clontech, Palo Alto, CA). After cloning and sequencing, nested primers specific for the 5' and 3' ends of the gene were synthesized. Another cycle of PCR was performed using Marathon Ready liver cDNA as a template and the full length cDNA was generated by PCR as follows: 1 cycle: 94°C 1min; 30 cycles: 94°C, 30 sec, 68°C, 7 min. The products of the reaction were subcloned into pCR2.1 (Invitrogen, Carlsbad, CA) and sequenced. The full-length cDNA of the 1.9 kb and 2.7 kb transcripts have been submitted to GenBank under accession

numbers AF208108 and AF208109, respectively.

Human cDNAs covering the coding region of the *EVI27* gene were synthesized in a similar manner as described above. Briefly, a human EST (accession number T96740), with high
5 homology to the mouse *EVI27* cDNA, was used to design nested oligonucleotide primers at the 5' and 3' end of the EST. 5' and 3' RACE products from Marathon Ready fetal liver cDNA (Clontech, Palo Alto, CA) were synthesized as described above. The 1.9 kb and 2.9 kb human *EVI27* cDNA nucleotide sequences are
10 deposited in GenBank under accession numbers AF208110 and AF208111.

EXAMPLE 8

Transfection and Stable Cell Lines

15 The 1.9 kb mouse *Evi27* cDNA was cloned into the *NotI-HindIII* site of the pcDNA3.1(-) eukaryotic expression vector and transformed into DH5 α bacteria. 10 μ g of recombinant plasmid DNA was resuspended at 1 μ g/ μ l in sterile TE. Cell lines 32Dcl3, M1, WEHI3B, and NIH3T3 were grown in appropriate
20 media maintaining cell viability at 90%. Cells were harvested and washed 3X in FBS-free DMEM media. Cells were resuspended at 5 x 10⁶ cells per ml in FBS-free DMEM. One ml of the cell suspension was mixed with DEAE dextran at 0.1 μ g/ml and 10 μ g of plasmid DNA in an 0.4 cm electrode gap electroporation
25 cuvette and chilled on ice for 10 minutes and Electroporated at 270 V/975 μ F in a Gene Pulser II (Bio-Rad, Hercules, CA). The cuvette was chilled on ice for 10 minutes then the cells were transferred into 10 ml of fresh media and grown for 72 hrs.

G418 selection was carried out in 2 mg/ml for 2 weeks. Limiting dilution cloning in 96 well culture dishes was performed at 1 mg/ml G418 on all cell lines with the exception of NIH3T3. Stable transfectants were cloned and tested for expression of the
5 transgene by Northern blot hybridization.

EXAMPLE 9

Southern and Northern Blot Probes

The β -actin probe was a 2.0 kb cDNA (Clontech, Palo Alto, CA). Probes pAKV5 and pEco have been described (Nakamura et al., 1996). p57BXH2A was a 700 bp PCR fragment generated from the flanking DNA from clone p57BXH2. Probe p57ET47 was a 103 bp exon trapped from p57 P1. The human intron probe was synthesized by PCR from genomic DNA. Human
10 ESTs were derived from IMAGE consortium clones. Full length 1.9 kb human and mouse *Evi27* cDNAs were used in all probes were labeled with [α^{32} P]-dCTP using the Prime It II labeling kit (Stratagene, La Jolla, CA).
15

EXAMPLE 10

DNA Sequencing

DNA sequencing was performed using the PRISM Ready Reaction DyeDeoxy Terminator Cycle Sequencing Kit (Perkin Elmer) on the ABI Model 373A DNA Sequencer (Applied Biosystems). Sequence primers were either the T3, T7
25 sequencing primers or synthetic oligomers derived from previously determined sequence.

EXAMPLE 11

Immunohistochemistry

Cells were harvested and washed once in PBS and resuspended at $1-2 \times 10^6$ cells per ml in PBS. 0.5 ml of the cell suspension was subjected to cytocentrifugation at $1200 \times g$ for 5-10 minutes and air dried for 15-20 minutes, rinsed in PBS and drained well but not allowed to dry. Slides were fixed in fresh 4% paraformaldehyde for 10 minutes followed by two washes in PBS.

Cells were permeabilized by incubation in acetone for 30 seconds at RT. Slides were rinsed in PBS with four changes over 5 minutes. Cells were blocked with 5% goat serum in PBS for 30 minutes at RT. The polyclonal anti-peptide EVI27 antibody, 2954, was affinity purified using the Sulfa-Link kit according to manufacturer's instructions (Pierce, Rockford, IL). Antibodies were diluted in 1 M HEPES buffer containing 0.15 N NaCl. Sixty microliters of the primary antibody solution was added to slides and incubated for 30 minutes in a humidified chamber at RT. The primary antibody was blocked with 100 μg of peptide by incubating for 1 hour at RT. This solution was then added to the slides as above. After hybridization the slides were washed three times for 5 minutes in 1XPBD (Oncor, Gaithersburg, MD). The secondary antibody, a rhodamine conjugated goat-anti-rabbit antibody (Pierce, Rockford, IL) was diluted in the same buffer as the primary antibody and the solution was added to slides for 30 minutes. Slides were washed 3X in PBS for 5 minutes each wash. The nuclei were counterstained with DAPI at 1/40 dilution in

antifade (Oncor, Gaithersburg, MD), coverslips added and viewed with an Olympus BX60 epifluoresence microscope.

EXAMPLE 12

5 Western Blotting

Cells were harvested at logarithmic growth stage and washed 2X in 1XPBS. Cells were resuspended in extraction buffer (1x PBS (pH 7.2), 10 µg/ml Aprotinin (0.1 unit/ml), 10 µg/ml Leupeptin, 1mM PMSF) at 10^6 cells/100 µl. Protein was extracted
10 by the freeze-thaw method. The solid phase was removed by centrifugation at 12,000g x 10 minutes at 4°C. The aqueous phase was collected concentrated and quantified using the BCA assay kit (Pierce, Rockford, IL). Protein separating gels were precast (Novex, San Diego, CA). Protein (50-100 µg) was mixed
15 1:1 with sample buffer (2X) (Novex, San Diego, CA), denatured for 15 minutes, cooled to RT and loaded. Gels were run for 2-3 hours at 200V with cooling (60W initially) in 1X running buffer (25 mM Tris, 192 mM glycine, 0.1% SDS at pH 8.3). Protein was transferred to Hybond-C super membrane (Amersham
20 Pharmacia Biotech, Piscataway, NJ) using an electrophoretic blotting system (C.B.S. Scientific Co. Del Mar, CA) in 1X transfer buffer (48mM Tris, 39 mM glycine, 20% methanol) at 100V for 5-6 hrs. The western blot was processed as described (WesternBreeze kit, Novex, San Diego, CA).

25

EXAMPLE 13

Computer DNA and Protein Sequence Analysis

Sequence homology searches were conducted at the

protein level using the National Center for Biotechnology Information and the BLAST network service. The DNA and protein sequence alignments and protein motif sequence searches were performed using MacVector (Oxford Molecular Group, Beaverton, OR) and pSORT (Nakai and Kanehisa, 1992).

EXAMPLE 14

Cloning and Characterization of Genomic Sequences from the *Evi27* locus

Genomic restriction analysis of DNA from two BXH2 leukemias with proviral integrations at *Evi27* (N57 [B160] and 15-38374) showed that the integrations in the two leukemias are located about 1.5 kb apart and are oriented in the same transcriptional direction (Figure 1). Lambda and P1 clones from the region were then isolated, restriction mapped and exon trapped. Four exon-trapped products were identified. BLAST searches showed that two of the exons were derived from the mouse choline dehydrogenase (*Chdh*) gene, while one exon was derived from a gene with high homology to the yeast actin-like protein, ARP8 (accession number S67026). The *Chdh* gene was subsequently positioned ~10 kb to the left of the proviral integration sites and in the opposite transcriptional direction (Figure 1). Northern analysis showed that *Chdh* expression is restricted to liver and is not activated in the B160 cell line (a cell line derived from the N57 leukemia that carries a proviral integration at *Evi27*). The ARP8 homologue was mapped ~15-20 kb to the right of the proviral integration sites. Northern analysis showed that this gene is ubiquitously expressed and is not

upregulated in B160 leukemia cells. On this basis, *Chdh* and *ARP8* were excluded as candidate disease genes.

The fourth exon (p57ET47) showed no significant homology to any sequences in GeneBank. Hybridization studies
5 showed that this exon was contained within lambda clone p57λ129 (Figure 1), indicating that it was located within 5-20 kb of the proviral integration sites at *Evi27*. DNA sequence analysis of p57λ129 identified a 103 bp fragment that was homologous to a human fetal liver cDNA EST (accession number T96740). This
10 EST sequence was localized 1240 bp upstream of p57ET47 and was found to be oriented in the same transcriptional direction, suggesting that both exons might be derived from the same gene.

Northern blot analysis of mouse tissues using both exons as probes confirmed this prediction and showed that the
15 *Evi27* hybridization pattern is complex. Six different *Evi27* transcripts of approximately 4.4, 4.2, 2.3, 1.9, 1.3, and 1.1kb in size, could be detected on northern blots (Figure 2). In adult tissues the expression was seen in liver and testes where the 2.3 and 1.9 kb transcripts predominated. Expression of the 4.2
20 transcript was also seen in liver and testes. In addition, low levels of the 2.3 and 1.9kb transcripts were seen in kidney (Figure 2A). With long exposure, the lung showed expression of a 2.1kb transcript, while the heart showed expression of the 1.3kb transcript. No expression was seen in skeletal muscle, brain, or
25 spleen.

In addition to adult tissues, *Evi27* expression was also detected in some hematopoietic cell lines. These cell lines include the EL-4 T-cell line, which express the 4.2 and 1.3 kb

transcripts, the WEHI231 B-cell line which expresses the 4.2, 1.3 and 1.1 kb transcripts, and the R1.1 T-cell line, which expresses the 1.9 kb transcript. In contrast, little or no expression was detected in the 32D, WEHI 3B, and M1 myeloid cell lines, the
5 HYB.548 B-cell line, or the P815 mast cell line (Figure 2C).

Evi27 expression could also be detected in BXH2 myeloid cell lines whether or not they carry a proviral integration at *Evi27*. Expression of the 4.4, 4.2, 1.9 and 1.1kb transcripts was variable among the different lines (Figure 2C), but the highest
10 expression was seen in the B160 cell line, which contains a viral integration at *Evi27*. In this cell line the 4.4 kb transcript predominated. These results suggest that viral integration at *Evi27* results in increased *Evi27* expression.

In humans, two *EVI27* transcripts, 1.9 kb and 2.7 kb in
15 size, were detected (Figure 3). Human *EVI27* expression is therefore considerably less complex than in mouse. The highest *EVI27* expression was seen in kidney (Figure 3A), while in the mouse, *Evi27* was expressed at low levels in the kidney. Moderate expression was also observed in the brain, liver, and testes with
20 low to undetectable expression in lung. No *EVI27* expression was seen in heart, placenta, and skeletal muscle (Figure 3A) or in immune tissues such as spleen, lymph nodes, thymus, peripheral blood lymphocytes, or bone marrow (Figure 3B). High *EVI27* expression was, however, observed in the fetal liver.

25 Within cell lines, moderate *EVI27* expression was seen in the cervical carcinoma HeLa, chronic myelogenous leukemia, K562, and colon adenocarcinoma, SW480 cell lines (Figure 3C), but not in the myeloid leukemia HL-60, acute lymphoblastic T-cell

leukemia MOLT-4, Burkitt's lymphoma Raji, lung carcinoma A549 and the melanoma G361 cell lines.

EXAMPLE 15

5 The Gene Whose Expression is Upregulated by Viral integration at Evi27 encodes a Novel IL-17 Receptor-Related Protein

The complete coding region of human gene was obtained by 3', and then 5' RACE. The primary 5' RACE product using primers from the 3' end of the gene was ~1.9 kb in length, 10 consistent with the size of the major human transcript. A larger minor species was also produced in the 5' RACE reaction. Sequence analysis showed that it contained an additional 950 bp of sequence that was not present in the 1.9 kb product. PCR and genomic sequence analysis showed that this extra sequence was 15 an unspliced intron. Northern analysis showed that the 2.7 kb human transcript contains this unspliced intron and sequence analysis showed that this unspliced intron introduces multiple stop codons into the open reading frame of the protein.

The sequence of the full length 1.9 kb and 2.7 kb 20 transcripts has been deposited in GenBank with accession numbers AF208110 and AF208111, respectively; SEQ. ID NO: 1 and SEQ. ID NO: 2 respectively in the present invention. The predicted uninterrupted protein is 502 amino acids (aa) and has a predicted molecular weight of 56 kD (Figure 4). pSORT analysis 25 identified a potential cleavable signal peptide at the N terminus of the protein. A putative transmembrane domain starts at residue 293 and ends at residue 309. The cytoplasmic tail is predicted to extend from residue 310 to the C terminus. pSORT analysis also

indicated that the protein has type 1a topology. A string of 39 amino acids derived from intronic sequences would be inserted at amino acid 250 in the 2.7 kb transcript. This would result in the synthesis of a 288 amino acids protein with a predicted molecular weight of 31 kD. The transmembrane and cytoplasmic portions of the 502 amino acids isoform are missing in this truncated form of the protein.

Multiple 5' RACE products were also obtained in mouse. Cloning and sequence analysis of these RACE products showed that they can be created by the inclusion of at least two unspliced introns. As in human, these intron sequences introduce multiple stop codons into the open reading frame of the protein. The sequence of the full length 1.9 kb and 2.5 kb mouse transcripts has been deposited in GenBank with accession numbers AF208108 and AF208109, respectively; SEQ. ID NO: 3 and SEQ. ID NO: 4 respectively in the present invention. The mouse 4.4, 4.2, 2.4 and 1.3 kb transcripts have not been cloned and sequenced.

Genomic sequence analysis indicates that the mouse gene consists of 11 exons and spans ~13 kb of genomic DNA (Figure 5). It is interesting to note that Evi27 cDNAs specifically lacking exon 10 have been cloned from mouse AML cell lines. The gene is transcribed in the same orientation as the integrated proviruses, which lie in a CpG rich region located approximately 6 kb 5' from the first known exon of the gene (Figure 1). The predicted open reading frame of the 1.9 kb transcript is 499 amino acids with a predicted molecular weight of 56 kD (Figure 4). pSORT analysis identified a potential cleavable signal peptide

in the N terminus of the protein, just like in the human protein. A putative transmembrane domain starts at amino acid 290 and ends at amino acid 293. The transmembrane domain is located in exon 12 of the gene and is spliced out in variant transcripts
5 sequenced from the WEHI-231 B-cell line. These variant transcripts splice out an exon containing the entire putative transmembrane domain. The lack of this exon does not disrupt the predicted open reading frame of the protein.

In the 2.5 kb transcript, a string of 57 novel amino
10 acids, derived from intron 7, is added at amino acid 162 creating a protein with an open reading frame of 218 amino acids and a predicted molecular weight of 24 kD. Like the shorter human isoform, this isoform would lack the transmembrane and cytoplasmic domains of the protein.

15 The full-length human and mouse proteins are 83% similar and 76% identical (Figure 4). The extracellular domain of both proteins contains three conserved N-linked glycosylation and a GSK3 phosphorylation site. A single conserved GSK3 phosphorylation site is also present in the cytoplasmic domain.

20 Amino acid sequence comparisons showed that *Evi27* has significant homology throughout its coding region to the human and mouse *IL-17 receptor (IL-17R)* (E value = 2.5×10^{-29}) (Figure 6). The position of the transmembrane domain with respect to the amino terminus is essentially the same in the two
25 proteins. However, the cytoplasmic tail of the *IL-17R* is nearly 304 amino acids longer than that of the *Evi27*. Although *Evi27* is predicted to encode both membrane bound and soluble forms, *IL-17R* is not known to encode a soluble form.

EXAMPLE 16

The Mouse and Human Genes map in Syntenic Regions

The human *EVI27* gene was mapped to chromosome 3p21 by fluorescence *in situ* hybridization of high-resolution G-banded chromosomes (Figure 7). These results are consistent with the mouse mapping data, which localize the mouse gene to chromosome 14 in a region of human 3p21 homology (Copeland et al., 1995). They also confirm that the human gene maps in a region of the human genome that is frequently rearranged in human myeloid leukemia.

EXAMPLE 17

Expression and Subcellular Localization of Evi27

The expression and subcellular localization of the mouse protein was examined using a polyclonal antibody raised against a peptide derived from the C-terminus of Evi27. Antibody specificity was confirmed by immunofluorescence staining of 32D cells, which express undetectable levels of *Evi27* mRNA (Figure 8A), and 32D cells stably transfected with a 2.7 kb *Evi27* cDNA clone. Western blot analysis showed that whereas the parental 32D cells show weak expression of the expected 55 kD full length protein, the transfected line expresses approximately 3-5 fold higher levels of this isoform (Figure 8B). Immunofluorescence analysis showed weak staining in the parental 32D cells whereas the transfected cells show very abundant staining on the surface with a capping pattern (Figure 8C, left panels). Surface expression of Evi27 was further shown by flow activated cell sorting analysis (Figure 8D). Clear cell surface staining was

observed after analysis of live cells or fixed, but not permeabilized, transfected cells (data not shown). In all experiments the staining could be quenched by preincubation of the antisera with Evi27 peptide (Figure 8C, right panels and Figure 8D), thus indicating the antiserum is specific for Evi27.

Murine T cell lymphoma EL-4 cells also express Evi27 on cell surface and the cell surface capping possibly indicates polydimerization and receptor activation of Evi27 (Figure 9). In contrast, Evi27 shows different localization in murine AML cells. Myeloid leukemia cells B187 and B160 both show diffuse speckled staining pattern in the cytoplasm, whereas the B190 cells show a capping pattern similar to the T cell lymphoma EL-4 (Figures 10-11). Moreover, abundant Evi27 protein was found in the cytoplasm, but not on the cell surface, of the B160 cells (Figure 11).

EXAMPLE 18

Western blot analysis of Evi27 Protein Expression

Western blot analysis of a number of leukemia cell lines confirmed the Northern results and showed that many different Evi27 isoforms are expressed at the protein level (Figure 12). A 56 kD protein, which is the predicted size of the full length protein containing the transmembrane and cytoplasmic domains, was expressed in most samples tested (Figure 12A). The EL4 and R1.1 T-cell lines and the WEHI 231 B-cell line also express a 24 kD protein, which is the predicted size of the truncated protein resulting from intron incorporation and an 85 kD protein (Figure 12A). This protein is also expressed at low levels in the

B190 cell line but not in the other the BXH2 leukemia cell lines tested.

Several cytokine receptors also produce soluble forms of the receptor by proteolytic cleavage of the membrane bound
5 receptor. The cleavage sites are located near the plasma membrane in the extracellular domain of the receptor. Proteolytic cleavage of the Evi27 membrane receptor in this region is predicted to produce a 33 kD protein. A band of this size is also present in most samples tested (Figure 12A). Several
10 unexpected sized Evi27 bands were also detected on the Western blot. These protein bands can all be competed away by the addition of Evi27 peptide (Figure 12B), indicating that they contain authentic Evi27 protein sequences.

The origin of these protein bands is unknown but
15 could result from the translation of Evi27 mRNA transcripts that have yet to be sequenced or may be indicative of homo- or heteropolymer formation. The expression of these protein bands is sometimes variable. For example, the 75 kD band is expressed at low or undetectable levels in the B160 and B187 cell lines. The
20 B193 cell lines does not express the 150 kD and 127 kD bands, but does expresses high levels of the 75 and 33 kD bands. The reason for this is not clear, but may reflect the differentiation state of the cell line. Consistent with this hypothesis, T- and B-cell lines express Evi27 protein bands that are not expressed in
25 BXH2 myeloid cell lines (i.e., 24 and 85 kD bands).

Evi27 expression was also examined in human hematopoietic cells. Western blot analysis showed that MEG01 (megakaryocytic leukemia), K562 (chronic myelogenous

leukemia) and HEL (erythrocytic leukemia) cell lines express the highest levels of Evi27 protein (Figure 13). Flow activated cell sorting analysis indicated that there is high level of surface expression on THP-1 (monocytic leukemia) and K562 cells but
5 not on HL-60 (promyelocytic leukemia) cells (Figure 14).

Discussion

10 The studies described here show that retroviral integration at the *Evi27* locus in BXH2 murine myeloid leukemias results in the increased expression of a novel gene with homology to the *IL-17 receptor*. A number of cytokine receptors (i.e., *v-mpl* or *Tpor*, *Il6ra*, *Epor*, *Il9r*, *Csf1*, *Il2ra*, *Il4ra*, and *Ifngr2*) or
15 their ligands (i.e., *Il6*, *Csf1*, *Il3*) are reported targets of retroviral integration or transduction in mouse leukemia/lymphomas making cytokines/cytokine receptors one of the largest and most important classes of leukemia genes (Ymer et al., 1985; Wendling et al., 1986; Penciolelli et al., 1987; Gisselbrecht et al., 1987;
20 Sugita et al., 1990; Blankenstein et al., 1990; Souyri et al., 1990; Hino et al., 1991; Chretien et al., 1994; Flubacher et al., 1994; Li et al., 1999). The *Evi27*-encoded cytokine-related receptor is therefore an excellent candidate for a new leukemia disease gene.

The human *EVI27* gene was also cloned and mapped to
25 chromosome 3p14-p21, in a region syntenic to the mouse gene. The human protein is 76% identical to the mouse protein at the amino acid level. This is somewhat higher than the recently described homology between the mouse and human IL-17

receptors, which are 69% identical at the amino acid level (Yao et al., 1997). This conservation suggests that Evi27 receptor function is evolutionarily conserved between human and mouse.

Proviral integrations at *Evi27* are located about 6 kb
5 upstream from the *IL-17 receptor*-related gene and result in increased gene expression. Viral integration at *Evi27* may thus induce disease simply by upregulating receptor expression. However, given the complex expression pattern observed for *Evi27*, it is also possible the viral integration induces disease by
10 altering *Evi27* isoform expression. In the mouse, as many as six *Evi27* transcripts are detected on Northern blots. Two of the transcripts result from intron incorporation and are predicted to produce a truncated soluble form of the receptor lacking the transmembrane and cytoplasmic domains. Several cytokine
15 receptors are known to produce soluble as well as membrane bound forms and it is now well documented that the soluble receptors can have both positive and negative effects on ligand signaling (for review see Heaney and Golde, 1996). The origin of the other *Evi27* transcripts has not been determined. It is thus
20 possible that one or more of these uncharacterized messages encode sequences that are located upstream of *Evi27* proviral integration sites at *Evi27*. If this is the case, then *Evi27* proviral integrations may prevent these messages from being expressed and this may have important disease consequences.

25 *IL-17r /Evi27* are unusual in that they have no homology with any protein in public databases, including other cytokine receptor proteins, and they have no recognizable motifs associated with intracellular signalling (Yao et al., 1995b). Many

disease-related cytokine receptors involved in leukemogenesis contain kinase domains; however, it is not unprecedented to find receptors lacking kinase domains that are involved in disease. For example, the human *MAS1* oncogene, which encodes a functional angiotensin receptor, is a G-protein coupled receptor that functions through phosphatidylinositol 4,5-bisphosphate hydrolysis (Young et al., 1986; Jackson et al., 1988). *Evi27* may thus identify a new disease pathway.

At least seven different *Evi27* isoforms (i.e., 24, 33, 56, 47, 75, 127, and 150 kD in size) were also detected in BXH2 leukemia cell extracts by western analysis using an *Evi27*-specific antibody. All leukemia extracts expressed the 56 kD isoform (the membrane bound receptor) as well as the 33 kD isoform (a postulated proteolytic cleavage containing the extracellular ligand-binding domain of the receptor). Expression of the other isoforms was variable among the extracts. One of seven cell extracts (B190) expressed the 24 kD isoform (a putative truncated receptor resulting from intron incorporation), three extracts (B139, B160, and B187) express the 47 kD isoform, five extracts (B112, B132, B139, B190, B193) the 75 kD isoform, and six extracts (B112, B132, B139, B160, B187, B190) the 150 kD isoform. The origin of these larger isoforms (i.e., oligomerization) or the reason for their variable expression (i.e. differentiation state of the cell) is presently unclear.

Only one BXH2 leukemia cell extract, B160, analyzed on Western blots is known to contain a proviral integration at *Evi27*. The 56 kD isoform is overexpressed in B160 cells, consistent with Northern results showing that viral integration at

Evi27 results in increased *Evi27* expression. Surprisingly, the 56 kD isoform is also overexpressed in B187 cells, which are not known to carry a viral integration at *Evi27*. Perhaps, B187 cells harbor a viral integration at *Evi27* that maps outside the region examined on Southern blots or there are other mechanisms for upregulating expression of this isoform other than proviral integration. It is interesting to note that B187 cells, like B160 cells, also fail to express the 75 kD isoform. One intriguing possibility is that viral integration at *Evi27* blocks expression of this isoform.

Evi27 expression is restricted in the hematopoietic cells. Northern analysis showed that *Evi27* is expressed in T cells and ESTs homologous to the mouse gene have been identified in CD4⁺ T-cells. Likewise, the mouse preB-cell line WEHI231 expresses *Evi27* and ESTs homologous to *EVI27* have been identified in human germinal center B cells. No *Evi27* expression was seen in the late stage B-cell line 548, the Burkitt lymphoma cell line Raji, the murine erythroid line D1b, or the mast cell line P815. A complex pattern of expression was observed within the myeloid compartment. Variable expression was seen in BXH2 leukemia cells, while little or no *Evi27* expression was seen in the M1 and 32D murine myeloid leukemia cell lines or in the monocytic leukemia cell line WEHI-3B. *Evi27* is expressed, however, in the chronic human myelogenous leukemia cell line K562, but not in the promyelocytic leukemia cell line HL-60. These results suggest that *Evi27* expression may be tightly regulated during myeloid cell differentiation and imply that *Evi27* may have an important function in controlling the growth and/or

differentiation of hematopoietic cells. Proviral integration at *Evi27* may interfere with this function(s) and, in doing so, lead to myeloid disease.

The *Evi27* subcellular protein distribution was also
5 analyzed in various mouse hematopoietic cell lines. In the EL4 T-cell and WEHI231 B-cell lines, *Evi27* protein was largely found on the cell surface in a capping pattern. Interestingly, only one BXH2 leukemia cell line, B190, has a similar staining pattern. In the other BXH2 leukemia cell lines examined, including B160,
10 which has a proviral integration at *Evi27*, *Evi27* protein is distributed throughout the cytoplasmic where it exhibits a punctate staining pattern. Western blot analysis shows that cells exhibiting the ER/Golgi staining pattern uniquely express the 24 kD *Evi27* isoform. It is possible that the 24 kD isoform uniquely
15 localizes to the ER/Golgi and somehow prevents the other *Evi27* isoforms from entering the cytoplasm. Alternatively, it is possible that a cofactor or heterodimeric partner required for transport from the ER/Golgi complex is differentially expressed. This hypothesis is strengthened by the fact that the 32D cell line
20 transfected with the 56kD isoform still lacks the cytoplasmic distribution seen in B160 or B187. In support of this possible mechanism, the expression of beta-2-microglobulin is required for cell surface expression of MHC class I or class I-like molecules (Lamouse-Smith et al., 1993; Feder et al., 1998). Additionally, the
25 receptor-associated protein (RAP), a type of chaperone, is especially designed to assist in the biosynthesis and intracellular transport of endocytic receptors (Willnow, 1998) and band 3 (AE1 gene) plays a chaperone-like role required for the

recruitment of Glycophorin A to the red blood cell plasma membrane (Hassoun et al., 1998). Finally, recent studies have shown that calnexin and Ig-alpha/Ig-beta interactions with membrane immunoglobulins are critical for the surface expression of the B cell antigen receptor of the IgM and IgD classes (Wu et al., 1997).

IL-17 (CTLA8) is a homodimeric cytokine of about 32 kD expressed exclusively from human memory T cells or mouse alpha beta TCR⁺CD4⁺CD8⁻ thymocytes (Rouvier et al., 1993; Yao et al 1995a; Yao et al., 1995b; Kennedy et al., 1996). In contrast with the tightly controlled expression of the ligand, the IL-17 receptor is ubiquitously distributed but more abundant in spleen and kidney (Yao et al., 1995b). Although devoid of direct effects on cells of hematopoietic origin, IL-17 induces the secretion of IL-6, IL-8, PGE2, MCP1 and G-CSF by adherent cells like fibroblasts, keratinocytes, epithelial and endothelial cells (Yao et al., 1995a; Yao et al., 1995b; Fossiez et al., 1996). When cultured in the presence of IL-17, fibroblasts can sustain the proliferation of CD34⁺ human progenitors and their preferential differentiation into neutrophils (Fossiez et al., 1996). Adenovirus-mediated transfer of murine IL-17 cDNA into liver has also been shown to induce a transient, but dramatic granulopoiesis *in vivo*, except in IL-6-deficient mice (Schwarzenberger et al., 1998).

The Evi27 protein is a member of the IL-17 receptor family. Two new members of the IL-17 cytokine family have recently been identified and these studies have demonstrated that IL-17B and IL-17C do not bind the IL-17 receptor extracellular domain, indicating that IL-17B and IL-17C are likely to bind

unique, yet uncharacterized receptors (Li et al., 2000). In a survey of cytokine induction, IL-17B and IL-17C stimulate the release of TNF- α and IL-1 β from the monocytic leukemia cell line THP-1 (Li et al., 2000). RT-PCR analysis has shown that IL-17B is
5 expressed by both the B160 leukemia and the stromal cell on which it depends. It is speculated that Evi27 may represent the receptor for IL-17B or IL017C. Based on these data a model for Evi27 in myeloid leukemia development is proposed. Terminal differentiation of myelomonocytic precursor cells likely result in
10 the down regulation of Evi27 expression. However, proviral insertions at Evi27 result in constitutive expression of the receptor. Binding of IL-17B/C to the Evi27 receptor would trigger the release of TNF- α and the IL-1 β by the leukemic cells. The TNF- α and IL-1 β would in turn provoke the production of multilineage
15 hematopoietic growth factors, adhesion molecules, and inflammatory cytokines by stromal cells (Bagby, 1994). These stromal cell derived factors then support the growth and survival of the leukemia cell and may account for the absolute dependence of the B160 leukemia on the stromal feeder layer for
20 growth and survival.

The human homologue of *Evi27* may also be involved in human disease. *EVI27* maps to chromosome 3p21, a region consistently deleted in a variety of human cancers. Loss of 3p heterozygosity is also frequently observed in renal cell
25 carcinoma, lung cancer and breast cancer and analysis of 3p allele loss in renal cell cancer has localized a candidate tumor suppressor gene to 3p21 (van den Berg et al., 1997). These results are consistent with the hypothesis that chromosome 3p

encodes a number of tumor suppressor genes. Given that *EVI27* maps to 3p21 and is expressed at high levels in normal human kidney, it will be interesting to determine whether this gene is affected by 3p21 mutations in renal cell carcinoma. *EVI27* also
5 may be a human myeloid leukemia disease gene. Recurrent treatment-related chromosome 3p21 breaks are frequently observed in myelodysplastic syndrome and acute myeloid leukemia patients (Shi et al., 1996), and 3p21 is the most frequently deleted region in human CML (Johansson et al., 1997).
10 Future studies will be aimed at determining whether *EVI27* is a human myeloid leukemia disease gene.

An Evi27 ligand, IL-17E, that induces activation of NF- κ B and stimulates production of the proinflammatory chemokine IL-8 was identified recently. Together with the restricted
15 expression of Evi27 disclosed herein, it is likely that Evi27 mediates the secretion of proinflammatory cytokines such as IL-8 and plays important role in the developmental and/or disease processes of hematopoietic cells. Hence, modulating the expression of Evi27 at the RNA or protein level may be exploited
20 for use in the treatment of diseases such as cancer or autoimmune diseases. A number of methods can be used to modulate Evi27 expression, e.g. by anti-sense oligonucleotides, small molecules that bind to the receptor, modified IL-17E ligand that may stimulate or repress the activity of the Evi27 receptor,
25 and the use of soluble form of the Evi27 receptor.

The following references were cited herein:

Bagby (1994). *The Molecular Basis of Blood Diseases*.

- Stamatoyannopoulos G, Nienhuis A.W., Majerus P.W. and Varmus H. (eds). W.B. Saunders Press: Philadelphia, p. 71-106.
- Baumbach et al., (1988). *J. Virol.*, **62**, 3151-5
- Bedigian et al., (1984). *J. Virol.*, **51**, 586-594.
- 5 Benton et al., (1977). *Science*, **196**, 180-2.
- Blankenstein et al., (1990). *J. Exp. Med.*, **171**, 965-70
- Buchburg et al., (1990) *Mol. Cell. Biol.*, **10**, 4658-4666.
- Chretien et al., (1994). *Blood*, **83**, 1813-1821
- Copeland et al., (1993). *Science*, **262**, 57-66.
- 10 Copeland NG, Jenkins NA 1999. Myeloid leukemia: disease genes and mouse models. In Animal Models of Cancer Predisposition Syndromes (eds Hiai, H. & Hino, O.) 53-63 (Karger, Basel, 1999).
- Feder et al., (1998). *Proc Natl Acad Sci U S A.*, **95**, 1472-7.
- Flubacher et al., (1994) *J. Virol.*, **68**, 7709-16
- 15 Fossiez et al., (1996). *J. Exp. Med.*, **183**, 2593-603.
- Gilbert et al., (1993). *J. Virol.*, **67**, 2083-90.
- Hassoun et al., (1998). *Blood* **91**, 2146-51.
- Heaney et al., (1996). *Blood*, **87**, 847-57
- Hino et al., (1991). *Mol. Cell. Biol.*, **11**, 5527-33.
- 20 Jackson et al., (1988). *Nature* **335**, 437-40
- Jenkins et al., (1982). *J. Virol.*, **42**, 379-388.
- Johansson et al., (1997). *Leukemia*, **11**, 1207-13.
- Kennedy et al., (1996) *J. Interferon Cytokine Res.*, **16**, 611-617.
- Lamouse-Smith et al., (1993). *J. Immunol.*, **15**, 6283-90.
- 25 Largaespada et al., (1995). *J. Virol.*, **69**, 5095-5102.
- Li et al., (1999). *Nat. Genet.*, **23**, 348-353.
- Li et al., (2000). *Proc. Natl. Acad. Sci. USA*, **97**, 773-778.
- Liao et al., (1997). *Oncogene*, **14**, 1023-9.

- Look AT. (1997). *Science*, **278**, 1059-64.
- Lupas, et al., (1991). *Science*, **252**, 1162-1164.
- Morishita et al., (1988). *Cell*, **54**, 831-40
- Nakai et al., (1992). *Genomics*, **14**, 897-911.
- 5 Nakamura, et al., (1996a) *Nat. Genet.* **12**, 149-153.
- Nakamura, et al., (1996b). *Nat. Genet.*, **12**, 154-158.
- Ogawa et al., (1996). *Oncogene*, **13**, 183-91.
- Roberts et al., (1998). *Hum. Mol. Genet.*, **7**, 1169-78
- Rouvier et al., (1993). *J Immunol* **150**, 5445-56.
- 10 Sambrook et al., *Molecular Cloning, A Laboratory Manual 2nd edn.* (Cold Spring Harbor, Laboratory Press New York, 1989).
- Schwarzenberger et al., (1998). *J. Immunol.*, **161**, 6383-9.
- Shaughnessy et al., (1999). *Oncogene*, **18**, 2069-84.
- Shi et al., (1996). *Cytogenet. Cell Genet.*, **74**, 295-9
- 15 Sugita et al., T. (1990). *J. Exp. Med.*, **171**, 2001-9.
- Viskochil et al., (1990). *Cell*, **62**, 187-92.
- van den Berg et al., (1997) *Genes Chromosomes Cancer*, **19**, 228-32.
- Willnow TE. (1998). *Biol. Chem.*, **379**, 1025-31.
- 20 Wu Y, Pun C, Hozumi N. (1997). *Immunol.*, **158**, 2762-70.
- Yao et al., (1995a). *Immunity*, **3**, 811-821.
- Yao et al., (1995b). *J. Immunol.*, **155**, 5483-6
- Yao et al., (1997). *Cytokine*, **9**, 794-800
- Young et al., (1986). *Cell*, **45**, 711-9.
- 25 Ymer et al., (1985). *Nature*, **317**, 255-8.

Any patents or publications mentioned in this specification are indicative of the levels of those skilled in the art to which the invention pertains. Further, these patents and

publications are incorporated by reference herein to the same extent as if each individual publication was specifically and individually indicated to be incorporated by reference.

One skilled in the art will appreciate readily that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those objects, ends and advantages inherent herein. The present examples, along with the methods, procedures, treatments, molecules, and specific compounds described herein are presently representative of preferred embodiments, are exemplary, and are not intended as limitations on the scope of the invention. Changes therein and other uses will occur to those skilled in the art which are encompassed within the spirit of the invention as defined by the scope of the claims.

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule encoding an IL-17 receptor-related protein selected from the group consisting
5 of:

(a) an isolated nucleic acid molecule of SEQ. ID NO: 1, 2, 3 or 4 which encodes an IL-17 receptor-related protein;

(b) an isolated nucleic acid molecule which is complimentary and hybridizes to the nucleic acid molecules of
10 (a); and

(c) isolated nucleic acid molecule differing from the isolated nucleic acid molecules of (a) or (b) in codon sequence due to the degeneracy of the genetic code, and which encodes an IL-17 receptor-related protein.

15

2. A fragment of nucleic acid molecule of claim 1 that is at least 10 bases in length and which will selectively hybridize to nucleic acid molecule encoding an IL-17 receptor-related protein.

20

3. The nucleic acid molecule of claim 2, wherein said nucleic acid molecule is used as an anti-sense molecule to inhibit the expression of an IL-17 receptor-related protein.

25

4. The nucleic acid molecule of claim 2, wherein said nucleic acid molecule is used for chromosomal mapping or mutation analysis of gene encoding an IL-17 receptor-related protein.

5. A genomic DNA encoding an IL-17 receptor-related protein, wherein said genomic DNA hybridizes to nucleic acid molecule of claim 1.

5 6. A vector comprising a nucleic acid molecule selected from the group consisting of SEQ. ID NO: 1, 2, 3 and 4 adapted for expression in a cell and has regulatory elements necessary for expression of said nucleic acid molecule in the cell.

10 7. A cell transfected with the vector of claim 6, wherein said vector expresses an IL-17 receptor-related protein.

8. The cell of claim 7, wherein said cell is selected from group consisting of bacterial cells, mammalian cells, plant
15 cells and insect cells.

9. An IL-17 receptor-related protein encoded by nucleic acid molecule selected from the group consisting of SEQ ID No. 3 and 4, wherein said protein is about 24, 33, 56, 47, 75,
20 127, and 150 kD in size as detected by western blot analysis in BXH2 leukemia cell.

10. A peptide derived from the protein of claim 9,
25 wherein said peptide is at least 4 amino acids in length.

11. An antibody that binds specifically to the protein of claim 9.

12. An antibody that binds specifically to the peptide of claim 10.

13. An IL-17 receptor-related protein encoded by
5 nucleic acid molecule of claim 1, wherein said protein has the amino acid sequence selected from the group consisting of SEQ. ID NO: 5, 6, 7 and 8.

14. A peptide derived from the protein of claim 13,
10 wherein said peptide is at least 4 amino acids in length.

15. An antibody that binds specifically to the protein of claim 13.

16. An antibody that binds specifically to the
15 peptide of claim 14.

17. A method of stimulating the secretion of cytokines from a cell, comprising the step of:
20 binding a ligand to the IL-17 receptor-related protein, Evi27, wherein said binding will activate Evi27 and results in secretion of cytokines from said cell.

18. The method of claim 17, wherein said cytokines
25 are selected from the group consisting of IL-1, IL-8 and TNF- α .

19. The method of claim 17, wherein said cell is selected from the group consisting of hematopoietic cells,

leukemia cells and kidney cells.

20. A method of modulating the expression and activity of an IL-17 receptor-related protein, Evi27, comprising
5 the step of:

contacting a molecule to a cell, wherein the binding of said molecule to Evi27 mRNA or protein results in increased or decreased expression and activity of the Evi27 protein.

10

21. The method of claim 20, wherein said molecule is selected from the group consisting of anti-sense oligonucleotides, small molecule that binds to Evi27, modified IL-17E and soluble form of the Evi27 receptor.




Fig. 1





Fig. 2C

Fig. 2B

Fig. 2A






Fig. 4




Fig. 5

IL17R 1 MGAARSPPSAVPGPLLGLLLLLLVLLISLAALC.....RSAVPRPQLVCSQPGLNC 50
 EV127 I MSLVLLISLAALC.....RSAVPRPQLVCSQPGLNC 24

IL17R 51 TVKNSTCLDSDSWTHPRNLTPSSPKDLQIQLFHFAHTQQGDLPFVAHTIEWTL 100
 EV127 25 GSETGPSPE--WMLQHDLPQDLRLDLRVEPVTTSVATGDYSILMNVSWVL 72

IL17R 101 QTDASILYL EGAELSVL-QLN TNERLCVRFELSKLRHHHR.....RWRFT 145
 EV127 73 RADASIRLLKATKICVTKGKSNFQSYSCVRCNRYTEAFQTQTRPSGGKWTFS 122

IL17R 146 FSHFVVDPDQEVETVTHHLPKPIPDGDPNHQSKNF LVPDCEHARNKVTTTP 195
 EV127 123 YIGFPVVELNTVYFJGAIHNIPNANMNEDEGPSMSVNF TSPQC LDHINKYKKK 172

IL17R 196 CMS SGLWDPNITVETLEAHQLRVSTLWNESTHYQILLTSPFPHMENHSC 245
 EV127 173 CVKAGSLWDPNITACKKNEETVEVNF TTTPLQNRYMALIQ.....HST 215

IL17R 246 FEHMHIPAPRPEEFHQRSNVTLTLRN LKQCCRHQVQIQPFFSSCLNDCL 295
 EV127 216 IIGFSQVFEPHQK-KQTRASVVIPTVTDGDS-EGATVQLTPYFPTCGSDCI 262

IL17R 296 RHSA TVSCPEMPDTPETIPDYMPLWVYWFITGJSILLVGSVILLIVCMTW 345
 EV127 263 RHKGITV V LCPQTGVFPFLDNNKSKPGGLPLLLLSLLVA.....TW 303

IL17R 346 RL A G P G S E K Y S D D T K Y T D G L P A A D L I P P P L K P R K V W T I Y S A D H P L Y V D V V 395
 EV127 304 V L V A G I Y L M W R H E R I K K T S F S T T T L L P P . . . I K V L V V Y P S - E I C F H H T I J 348

IL17R 396 L K F A Q F L L T A C G T E V A L D L E E Q A I S E A G V M T W V G R Q K Q E M V E S N S K I I V 445
 EV127 349 C Y F T E F L Q N H C R S E V I L E K W Q K K I A E M G P V O W L A T Q K K . . . A A D K V V F 394

Fig. 6-1

446 495
 395 439
 IL17R L C S R G T R A K W Q A L L G R G A P V R L R C D H K P V G D L F T A A M N M I L P D F K R P A C
 EV127 I L S N D V N S V C D G T C G K S E G S P S E N S Q D L F P L A F N L F C S D L R S Q I H

 496 545
 440 478
 IL17R F G T Y V V C Y F S E V S C D G D V P D L F G A P R Y P L M D R F E E V Y F R I Q D L E M F Q P G
 EV127 L H K Y V V V Y F R E I D T K D D Y N A L S - V C P K Y H F M K D A T A F C A E

 546 595
 479 502
 IL17R R M H R V G E L S G D N Y L R S P P G R Q L R A A L D R F R D W Q V R C P D W F E C E N L Y S A D D
 EV127 L L H V K Q Q V S A G K R S Q A C H D G C C S L
 (SEQ ID No. 5)

 596 645
 IL17R Q D A P S L D E E V F E E P L L P P G T G I V K R A P L V R E P G S Q A C L A I D P L V G E E G G A

 646 695
 IL17R A V A K L E P H L Q P R G Q P A P Q P L H T L V L A A E E G A L V A A V E P G P L A D G A A V R L A


 696 745
 IL17R L A G E G E A C P L L G S P G A G R N S V L F L P V D P E D S P L G S S T P M A S P D L L P E D V R

 746 795
 IL17R E H L E G L M L S L F E Q S L S C Q A Q G G C S R P A M V L T D P H T P Y E E E Q R Q S V Q S D Q G

 796 845
 IL17R Y I S R S S P Q P P E G L T E M E E E E E E Q D P G K P A L P L S P E D L E S L R S L Q R Q L L F

 846
 IL17R R Q L Q K N S G W D T M G S E S E G P S A ⁸⁴⁶
 (SEQ ID No. 9)

Fig. 6-2






Fig. 7B




Fig. 8A

Fig. 8B




Fig. 8C




Fig. 8D




Fig. 9




Fig. 10





Fig. 11




Fig. 12B




Fig. 12A




Fig. 13




Fig. 14

SEQUENCE LISTING

<110> Shaughnessy, John D.
 <120> Evi27 Gene Sequence and Protein Encoded Thereby
 <130> D6138PCT
 <140>
 <141> 2001-02-02
 <150> US 60/180,374
 <151> 2000-02-04
 <160> 9
 <210> 1
 <211> 1827
 <212> DNA
 <213> *Homo sapiens*
 <220>
 <221> prim_transcript
 <223> cDNA of human Evi27

<400> 1
 cggcgatgtc gctcgtgctg ataagcctgg ccgcgctgtg caggagcgcc 50
 gtaccccgag agccgaccgt tcaatgtggc tctgaaactg ggccatctcc 100
 agagtggatg ctacaacatg atctaattccc cggagacttg agggacctcc 150
 gagtagaacc tgttacaact agtggtgcaa caggggacta ttcaattttg 200
 atgaatgtaa gctgggtact ccgggcagat gccagcatcc gcttggtgaa 250
 ggccaccaag atttgtgtga cgggcaaaag caacttccag tcctacagct 300
 gtgtgaggtg caattacaca gaggccttcc agactcagac cagaccctct 350
 ggtggtaaat ggacattttc ctatatcggc ttccctgtag agctgaacac 400
 agtctatttc attggggccc ataatatctc taatgcaa atgaatgaag 450
 atggcccttc catgtctgtg aatttcacct caccaggctg cctagaccac 500
 ataatgaaat ataaaaaaaa gtgtgtcaag gccggaagcc tgtgggatcc 550
 gaacatcact gcttgtaaga agaattgagga gacagtagaa gtgaacttca 600
 caaccactcc cctgggaaac agatacatgg ctcttatcca acacagcact 650
 atcatcgggt tttctcaggt gtttgagcca caccagaaga aacaaacgcg 700
 agcttcagtg gtgattccag tgactgggga tagtgaaggt gctacggtgc 750
 agctgactcc atattttcct acttggtggca gcgactgcat ccgacataaa 800
 ggaacagttg tgctctgccc acaaacaggc gtccctttcc ctctggataa 850
 caacaaaagc aagccgggag gctggctgcc tctcctcctg ctgtctctgc 900
 tgggtggccac atgggtgctg gtggcaggga tctatcta atgtggaggcac 950
 gaaaggatca agaagacttc cttttctacc accacactac tgccccccat 1000
 taaggttcct gtgggtttacc catctgaa atgtttccat cacacaattt 1050
 gttacttcac tgaatttcct caaaaccatt gcagaagtga gggtcatcctt 1100
 gaaaagtggc agaaaaagaa aatagcagag atgggtccag tgcagtggct 1150
 tgccactcaa aagaaggcag cagacaaagt cgtcttcctt ctttccaatg 1200
 acgtcaacag tgtgtgcgat ggtacctgtg gcaagagcga gggcagtcct 1250
 agtgagaact ctcaagacct cttccccctt gcctttaacc ttttctgcag 1300
 tgatctaaga agccagattc atctgcacaa atacgtggtg gtctacttta 1350
 gagagattga tacaaaagac gattacaatg ctctcagtg ctgccccaa 1400
 taccacttca tgaaggatgc cactgcttct tgtgcagaac ttctccatgt 1450
 caagcagcag gtgtcagcag gaaaaagatc acaagcctgc cagcatggct 1500
 gctgctcctt gtagccacc catgagaagc aagagacctt aaaggcttcc 1550

tatccccacca	attacagggga	aaaaacgtgt	gatgatcctg	aagcttacta	1600
tgcagcctac	aaacagcctt	agtaattaaa	acattttata	ccaataaaaat	1650
tttcaaataat	tactaactaa	tgtagcatta	actaacgatt	ggaaactaca	1700
tttacaactt	caaagctgtt	ttatacatag	aaatcaatta	cagctttaat	1750
tgaaaactgt	aaccattttg	ataatgcaac	aataaagcat	cttccaaaaa	1800
aaaaaaaaaa	aaaaaaaaaa	aaaaaaa			1827

<210> 2
 <211> 2856
 <212> DNA
 <213> *Homo sapiens*

<220>

<221> prim_transcript
 <223> cDNA of human Evi27

<400> 2					
cggcgatgtc	gctcgtgctg	ataagcctgg	ccgcgctgtg	caggagcgcc	50
gtaccccagag	agccgaccgt	tcaatgtggc	tctgaaactg	ggccatctcc	100
agagtggatg	ctacaacatg	atctaattccc	cggagacttg	agggacctcc	150
gagtagaacc	tgttacaact	agtgttgcaa	caggggacta	ttcaattttg	200
atgaatgtaa	gctgggtact	ccgggcagat	gccagcatcc	gcttggtgaa	250
ggccaccaag	atltgtgtga	cgggcaaaag	caacttccag	tcctacagct	300
gtgtgaggtg	caattacaca	gaggccttcc	agactcagac	cagaccctct	350
ggtggtaaat	ggacattttc	ctatatcggc	ttccctgtag	agctgaacac	400
agtctatttc	attggggccc	ataatattcc	taatgcaaat	atgaatgaag	450
atggcccttc	catgtctgtg	aatttcacct	caccaggctg	cctagaccac	500
ataatgaaat	ataaaaaaaaa	gtgtgtcaag	gccggaagcc	tgtgggatcc	550
gaacatcact	gcttgtaaga	agaatgagga	gacagtagaa	gtgaacttca	600
caaccactcc	cctgggaaac	agatacatgg	ctcttatcca	acacagcact	650
atcatcgggt	tttctcaggt	gtttgagcca	caccagaaga	aacaaaacgcg	700
agcttcagtg	gtgattccag	tgactgggga	tagtgaaggt	gctacgggtc	750
aggtaaagtt	cagtgaagctg	ctctggggag	ggaagggaca	tagaagactg	800
ttccatcatt	cattgctttt	aaggatgagt	tctctcttgt	caaatgcact	850
tctgccagca	gacaccagtt	aagtggcggt	catggggggt	ctttcgctgc	900
agctccacc	gtgctgaggt	caggaggccg	acgtggcagt	tgtggtcctt	950
tttgcttgta	ttaatggctg	ctgaccttcc	aaagcacttt	ttattttcat	1000
tttctgtcac	agacactcag	ggatagcagt	accattttac	ttccgcaagc	1050
ctttaactgc	aagatgaagc	tgcaaagggt	ttgaaatggg	aagggtttgag	1100
ttccaggcag	cgtatgaact	ctggagaggg	gctgccagtc	ctctctgggc	1150
cgcagcggac	ccagctggaa	cacaggaagt	tggagcagta	ggtgctcctt	1200
cacctctcag	tatgtctctt	tcaactctag	tttttgaagt	ggggacacag	1250
gaagtccagt	ggggacacag	ccactcccca	aagaataagg	aacttccatg	1300
cttcattccc	tggcataaaa	agtgnatcaa	cacaccagag	ggggcaggga	1350
ccagccaggg	tatgatgggt	actacccttt	tctggagaac	catagacttc	1400
ccttactaca	gggacttgca	tgtcctaaag	cactggctga	aggaagccaa	1450
gaggatcact	gctgctcctt	ttttgtagag	gaaatgtttg	tgtacgtggt	1500
aagatatgac	ctagcccttt	taggtaagcg	aactggatatg	ttagtaacgt	1550
gtacaaagtt	taggttcaga	ccccgggagt	cttgggcatg	tgggtctcgg	1600
gtcactgggt	ttgactttag	ggctttgtta	cagatgtgtg	accaagggga	1650
aaatgtgcat	gacaacacta	gaggttaggg	cgaagccaga	aagaagggaa	1700
gttttggctg	aagtaggagt	cttgggtgaga	ttttgctgtg	atgcatgggtg	1750
tgaactttct	gagcctcttg	tttttccctca	gctgactcca	tatttttcta	1800
cttgtggcag	cgactgcctc	cgacataaag	gaacagttgt	gctctgcccc	1850
caaacaggcg	tccctttccc	tctggataac	aacaaaagca	agccgggagag	1900

```

ctggetgcct ctcctcctgc tgtctctgct ggtggccaca tgggtgctgg 1950
tggcagggat ctatctaata tggaggcacg aaaggatcaa gaagacttcc 2000
ttttctacca ccacactact gccccccatt aagggttcttg tggtttacct 2050
atctgaaata tgtttccatc acacaatttg ttacttccact gaatttcttc 2100
aaaaccattg cagaagtgag gtcattcctg aaaagtggca gaaaaagaaa 2150
atagcagaga tgggtccagt gcagtggcct gccactcaaa agaaggcagc 2200
agacaaagtc gtcttccttc tttccaatga cgtcaacagt gtgtgcatg 2250
gtacctgtgg caagagcgag ggcagtccca gtgagaactc tcaagacctc 2300
ttcccccttg cctttaacct tttctgcagt gatctaagaa gccagattca 2350
tctgcacaaa tacgtgggtg tctacttttag agagattgat acaaaagacg 2400
attacaatgc tctcagtgtc tgccccaagt accacttcat gaaggatgcc 2450
actgctttct gtgcagaact tctccatgtc aagcagcagg tgtcagcagg 2500
aaaaagatca caagcctgcc acgatggctg ctgctccttg tagcccaccc 2550
atgagaagca agagacctta aaggcttcct atcccaccaa ttacagggaa 2600
aaaacgtgtg atgatcctga agcttactat gcagcctaca aacagcctta 2650
gtaattaaaa cattttatac caataaaatt ttcaaataat actaactaat 2700
gtagcatata ctaacgattg gaaactacat ttacaacttc aaagctgttt 2750
tatacataga aatcaattac agctttaatt gaaaactgta accattttga 2800
taatgcaaca ataaagcatc ttccaaaaaa aaaaaaaaaa aaaaaaaaaa 2850
aaaaaa

```

```

<210>      3
<211>      1963
<212>      DNA
<213>      Unknown

```

```

<220>

```

```

<221>      prim_transcript
<223>      cDNA of mouse Evi27

```

```

<400>      3
gtggccagtg gccgggccat gttgctagtg ttgctgatct tggctgcatc 50
gtgcaggagc gccctgcctc gagagccgac tattcagtgt ggctctgaga 100
cagggccatc tccagagtgg atgggtccaa acacactcac tccaggagac 150
ttgagggacc tccaagtgga actcgtcaag acaagtgtgg cagcagagga 200
gttttcaatt ttgatgaaca taagctggat actccgggca gacgccagca 250
tccgcttggt gaaggccacc aagatctgcg tgagtggcaa aaacaacatg 300
aattcatata gctgtgtgag gtgcaactac acagaggcct tccaaagcca 350
gaccagacct tccggcggca aatggacatt ctctatgta ggcttccctg 400
tgagctgag cactctctat ctcatcagcg ccataacat ccccaatgct 450
aatatgaatg aggacagccc ttctttgtct gtgaacttca cctcgccagg 500
ctgcctaaac cacgtaatga aatataaaaa gcagtgcact gaggcgggaa 550
gcctgtggga cccagacatc actgcttgta aaaagaacga gaagatgggt 600
gaagtgaatt tcacaaccaa tccccttgga aacagatata cgatttctcat 650
tcaacgggac acgacattgg ggttttctag agtgctggag aataaactga 700
tgaggacgtc tgtagccatc cgggtgactg aggagagtga aggtgcgggtg 750
gttcagctga ccccatattt acatacctgc ggcaatgact gcatccgacg 800
cgaagggaca gttgtgcttt gctcagagac aagtgtctcc atccctccag 850
atgacaacag acgcatgctg ggaggctggc tgctctctct cctgggtgctg 900
ctggtggctg tgtgggtgct ggcagctggg atctacctaa cttggaggca 950
aggaaggagc acgaagacgt cctttcctat ttccaccatg ctctgcccc 1000
tcattaaggt cctgggtggt tatccttctg agatatgttt ccatcacacc 1050
gtctgtcgct tcaactgact tcttcaaaac tactgcagaa gtgaggtcat 1100
ccttgaaaaa tggcagaaaa agaaaatcgc cgagatgggg ccggtacagt 1150
ggctgaccac tcagaagcaa gcggcagata aagtgggtctt ccttcttccc 1200

```

```

agtgacgtcc cgaccctttg tgacagtgcc tgtggccaca atgagggcag 1250
cgccagggag aactctcagg atctgttccc tcttgccctt aacctctttt 1300
gtagtgattt cagcagccag acgcatctgc acaaatacct ggtggtctat 1350
cttggggggag cagacctcaa aggcgactat aatgccctga gtgtctgccc 1400
ccaatatcat ctcatgaagg acgccacagc tttccacaca gaacttctca 1450
aggctacgca gagcatgtca gtgaagaaac gtcacaaagc ctgccatgat 1500
agctgttcac ccttgtagtc caccgggggg aatagagact ctgaagcctt 1550
cctactctcc cttccagtga caaatgctgt gtgacgactc tgaaatgtgt 1600
gggagaggct gtgtggagggt agtgctatgt acaaacttgc tttaaaactg 1650
gagtttgcaa agtcaacctg agcatacacg cctgaggcta gtcattggct 1700
ggatttatga agacaacaca gttacagaca ataatgagtg ggacctacat 1750
ttgggatata cccaaagctg ggtaatgatt atcactgaga accacgact 1800
ctggccatga agtaatacgg cacttccctg tcaggctgtc tgtcagggtt 1850
ggtctgtctt gcactgcccc tgctctatgc tgcacgtaga ccgttttgta 1900
acattttaat ctgttaatga ataatccgtt tgggaagctc tcaaaaaaaaa 1950
aaaaaaaaaa aaa 1963

```

```

<210>      4
<211>      2589
<212>      DNA
<213>      Unknown

```

```
<220>
```

```

<221>      prim_transcript
<223>      cDNA of mouse Evi27

```

```

<400>      4
gtggccagtg gccgggccat gttgctagtg ttgctgatct tggtctgcatc 50
gtgcaggagc gccctgcctc gagagccgac tattcagtgt ggctctgaga 100
cagggccatc tccagagtgg atggtccaac acacactcac tccaggagac 150
ttgagggacc tccaagtgga actcgtcaag acaagtgtgg cagcagagga 200
gttttcaatt ttgatgaaca taagctggat actccgggca gacgccagca 250
tccgcttggt gaaggccacc aagatctgcg tgagtggcaa aaacaacatg 300
aattcataca gctgtgtgag gtgcaactac acagaggcct tccaaagcca 350
gaccagacct tccggcggca aatggacatt ctctatgta ggcttccctg 400
tgagctgag cactctctat ctctcagcg ccataacat cccaatgct 450
aatatgaatg aggacagccc ttctttgtct gtgaacttca cctcgccagg 500
gtgcactcgt gaaaacacag aagtaacgtc cgggtgattt ccagcagcta 550
aacaccaggc tctccggatt tcagctcctt tcccattaca atttcctcct 600
gggccagagg actcagtcct tctgccacc cagcctctgg cgtcgctttt 650
tcatgacttt gtcaaactta cctagcttgt ttccattctg aaattgtctg 700
atgcttgctt cgtatgtaag ccggggatat gaggtttggg tatgaatccc 750
acagagggca ctgaattcct ctactatgg cctatctggg ctgtgtgaca 800
ttgttggtga gggtcgtgcc tactaggcat ctgggtatct accacctgga 850
gcttcattgtc tggagagggc agaacctata tgtattgtca gctctcactt 900
ttgtttccgt gtcacctcct ggagactgtt tttgataaag gttgtactta 950
aaggagatta cttaaagcct ccgtggaaga atggtttcct atttagatct 1000
gttgtctctc atatctgaag taagtgtgtg tgtgtgtgtt ttgtgtgtgt 1050
gtgtgtgtgt gtgtgtgtac tgggcaaagg gttatacctt tactcaaattg 1100
taacaacttt cattcacatt cccaggctgc ctaaaccacg taatgaaata 1150
taaaaagcag tgcactgagg cgggaagcct gtgggacca gacatcactg 1200
cttgtaaaaa gaacgagaag atggttgaag tgaatttcac aaccaatccc 1250
cttggaacaa gatacacgat tctcattcaa cgggacacga cattgggggtt 1300
ttctagagtg ctggagaata aactgatgag gacgtctgta gccatcccg 1350
tgactgagga gagtgaaggt gcggtggttc agctgacccc atatttacat 1400

```

```

acctgcgga atgactgcat ccgacgcgaa gggacagttg tgctttgctc 1450
agagacaagt gctcccatcc ctccagatga caacagacgc atgctgggag 1500
gctgggtgcc tctcttcctg gtgctgctgg tggctgtgtg ggtgctggca 1550
gctgggatct acctaaacttg gaggaagga aggagcacga agacgtcctt 1600
tcctatttcc accatgctcc tgcccctcat taaggctcctg gtggtttatc 1650
cttctgagat atgtttccat cacaccgtct gtcgcttcac tgactttctt 1700
caaaactact gcagaagtga ggtcatcctt gaaaaatggc agaaaaagaa 1750
aatcgccgag atggggccgg tacagtggct gaccactcag aagcaagcgg 1800
cagataaagt ggtcttcctt ctccccagtg acgtcccgc cctttgtgac 1850
agtgcctgtg gccacaatga gggcagcgcc agggagaact ctcaggatct 1900
gttccctctt gcctttaacc tctttttag tagtttcagc agccagacgc 1950
atctgcacaa atacctgggtg gtctatcttg ggggagcaga cctcaaaggc 2000
gactataatg ccctgagtgt ctgcccccaa tatcatctca tgaaggacgc 2050
cacagctttc cacacagaac ttctcaaggc tacgcagagc atgtcagtga 2100
agaaacgctc acaagcctgc catgatagct gttcaccctt gtagtccacc 2150
cgggggaata gagactctga agccttecta ctctcccttc cagtacaaa 2200
tgctgtgtga cgactctgaa atgtgtggga gaggtgtgtg ggaggtagt 2250
ctatgtacaa acctgttcta aaactggagt ttgcaaagtc aacctgagca 2300
tacacgcctg aggtagtca ttggctggat ttatgaagac aacacagtta 2350
cagacaataa tgagtgggac ctacatttgg gatataccca aagctgggta 2400
atgattatca ctgagaacca cgcactctgg ccatgaagta atacggcact 2450
tccctgtcag gctgtctgtc aggttgggtc tgtcttgac tgcccatgct 2500
ctatgctgca cgtagaccgt tttgtaacat tttaatctgt taatgaataa 2550
tccgtttggg aagctctcaa aaaaaaaaaa aaaaaaaaaa 2589

```

```

<210>      5
<211>      502
<212>      PRT
<213>      Homo sapiens

```

```

<220>

```

```

<221>      PEPTIDE
<223>      Human Evi27 protein

```

```

<400>      5
Met Ser Leu Val Leu Ile Ser Leu Ala Ala Leu Cys Arg Ser Ala
                    5                      10                      15
Val Pro Arg Glu Pro Thr Val Gln Cys Gly Ser Glu Thr Gly Pro
                    20                      25                      30
Ser Pro Glu Trp Met Leu Gln His Asp Leu Ile Pro Gly Asp Leu
                    35                      40                      45
Arg Asp Leu Arg Val Glu Pro Val Thr Thr Ser Val Ala Thr Gly
                    50                      55                      60
Asp Tyr Ser Ile Leu Met Asn Val Ser Trp Val Leu Arg Ala Asp
                    65                      70                      75
Ala Ser Ile Arg Leu Leu Lys Ala Thr Lys Ile Cys Val Thr Gly
                    80                      85                      90
Lys Ser Asn Phe Gln Ser Tyr Ser Cys Val Arg Cys Asn Tyr Thr
                    95                      100                     105
Glu Ala Phe Gln Thr Gln Thr Arg Pro Ser Gly Gly Lys Trp Thr
                    110                     115                     120
Phe Ser Tyr Ile Gly Phe Pro Val Glu Leu Asn Thr Val Tyr Phe
                    125                     130                     135
Ile Gly Ala His Asn Ile Pro Asn Ala Asn Met Asn Glu Asp Gly
                    140                     145                     150

```


Pro	Ser	Met	Ser	Val	Asn	Phe	Thr	Ser	Pro	Gly	Cys	Leu	Asp	His
				155					160					165
Ile	Met	Lys	Tyr	Lys	Lys	Lys	Cys	Val	Lys	Ala	Gly	Ser	Leu	Trp
				170					175					180
Asp	Pro	Asn	Ile	Thr	Ala	Cys	Lys	Lys	Asn	Glu	Glu	Thr	Val	Glu
				185					190					195
Val	Asn	Phe	Thr	Thr	Thr	Pro	Leu	Gly	Asn	Arg	Tyr	Met	Ala	Leu
				200					205					210
Ile	Gln	His	Ser	Thr	Ile	Ile	Gly	Phe	Ser	Gln	Val	Phe	Glu	Pro
				215					220					225
His	Gln	Lys	Lys	Gln	Thr	Arg	Ala	Ser	Val	Val	Ile	Pro	Val	Thr
				230					235					240
Gly	Asp	Ser	Glu	Gly	Ala	Thr	Val	Gln	Leu	Thr	Pro	Tyr	Phe	Pro
				245					250					255
Thr	Cys	Gly	Ser	Asp	Cys	Ile	Arg	His	Lys	Gly	Thr	Val	Val	Leu
				260					265					270
Cys	Pro	Gln	Thr	Gly	Val	Pro	Phe	Pro	Leu	Asp	Asn	Asn	Lys	Ser
				275					280					285
Lys	Pro	Gly	Gly	Trp	Leu	Pro	Leu	Leu	Leu	Leu	Ser	Leu	Leu	Val
				290					295					300
Ala	Thr	Trp	Val	Leu	Val	Ala	Gly	Ile	Tyr	Leu	Met	Trp	Arg	His
				305					310					315
Glu	Arg	Ile	Lys	Lys	Thr	Ser	Phe	Ser	Thr	Thr	Thr	Leu	Leu	Pro
				320					325					330
Pro	Ile	Lys	Val	Leu	Val	Val	Tyr	Pro	Ser	Glu	Ile	Cys	Phe	His
				335					340					345
His	Thr	Ile	Cys	Tyr	Phe	Thr	Glu	Phe	Leu	Gln	Asn	His	Cys	Arg
				350					355					360
Ser	Glu	Val	Ile	Leu	Glu	Lys	Trp	Gln	Lys	Lys	Lys	Ile	Ala	Glu
				365					370					375
Met	Gly	Pro	Val	Gln	Trp	Leu	Ala	Thr	Gln	Lys	Lys	Ala	Ala	Asp
				380					385					390
Lys	Val	Val	Phe	Leu	Leu	Ser	Asn	Asp	Val	Asn	Ser	Val	Cys	Asp
				395					400					405
Gly	Thr	Cys	Gly	Lys	Ser	Glu	Gly	Ser	Pro	Ser	Glu	Asn	Ser	Gln
				410					415					420
Asp	Leu	Phe	Pro	Leu	Ala	Phe	Asn	Leu	Phe	Cys	Ser	Asp	Leu	Arg
				425					430					435
Ser	Gln	Ile	His	Leu	His	Lys	Tyr	Val	Val	Val	Tyr	Phe	Arg	Glu
				440					445					450
Ile	Asp	Thr	Lys	Asp	Asp	Tyr	Asn	Ala	Leu	Ser	Val	Cys	Pro	Lys
				455					460					465
Tyr	His	Phe	Met	Lys	Asp	Ala	Thr	Ala	Phe	Cys	Ala	Glu	Leu	Leu
				470					475					480
His	Val	Lys	Gln	Gln	Val	Ser	Ala	Gly	Lys	Arg	Ser	Gln	Ala	Cys
				485					490					495
His	Asp	Gly	Cys	Cys	Ser	Leu								
				500										

<210> 6
 <211> 288
 <212> PRT
 <213> *Homo sapiens*
 <220>

<221> PEPTIDE
<223> Human Evi27 protein

<400> 6
Met Ser Leu Val Leu Ile Ser Leu Ala Ala Leu Cys Arg Ser Ala
5 10 15
Val Pro Arg Glu Pro Thr Val Gln Cys Gly Ser Glu Thr Gly Pro
20 25 30
Ser Pro Glu Trp Met Leu Gln His Asp Leu Ile Pro Gly Asp Leu
35 40 45
Arg Asp Leu Arg Val Glu Pro Val Thr Thr Ser Val Ala Thr Gly
50 55 60
Asp Tyr Ser Ile Leu Met Asn Val Ser Trp Val Leu Arg Ala Asp
65 70 75
Ala Ser Ile Arg Leu Leu Lys Ala Thr Lys Ile Cys Val Thr Gly
80 85 90
Lys Ser Asn Phe Gln Ser Tyr Ser Cys Val Arg Cys Asn Tyr Thr
95 100 105
Glu Ala Phe Gln Thr Gln Thr Arg Pro Ser Gly Gly Lys Trp Thr
110 115 120
Phe Ser Tyr Ile Gly Phe Pro Val Glu Leu Asn Thr Val Tyr Phe
125 130 135
Ile Gly Ala His Asn Ile Pro Asn Ala Asn Met Asn Glu Asp Gly
140 145 150
Pro Ser Met Ser Val Asn Phe Thr Ser Pro Gly Cys Leu Asp His
155 160 165
Ile Met Lys Tyr Lys Lys Lys Cys Val Lys Ala Gly Ser Leu Trp
170 175 180
Asp Pro Asn Ile Thr Ala Cys Lys Lys Asn Glu Glu Thr Val Glu
185 190 195
Val Asn Phe Thr Thr Thr Pro Leu Gly Asn Arg Tyr Met Ala Leu
200 205 210
Ile Gln His Ser Thr Ile Ile Gly Phe Ser Gln Val Phe Glu Pro
215 220 225
His Gln Lys Lys Gln Thr Arg Ala Ser Val Val Ile Pro Val Thr
230 235 240
Gly Asp Ser Glu Gly Ala Thr Val Gln Val Lys Phe Ser Glu Leu
245 250 255
Leu Trp Gly Gly Lys Gly His Arg Arg Leu Phe His His Ser Leu
260 265 270
Leu Leu Arg Met Ser Ser Leu Leu Ser Asn Ala Leu Leu Pro Ala
275 280 285
Asp Thr Ser
288

<210> 7
<211> 499
<212> PRT
<213> Unknown

<220>

<221> PEPTIDE
<223> Mouse Evi27 protein

<400> 7

Met	Leu	Leu	Val	Leu	Leu	Ile	Leu	Ala	Ala	Ser	Cys	Arg	Ser	Ala
				5					10					15
Leu	Pro	Arg	Glu	Pro	Thr	Ile	Gln	Cys	Gly	Ser	Glu	Thr	Gly	Pro
				20					25					30
Ser	Pro	Glu	Trp	Met	Val	Gln	His	Thr	Leu	Thr	Pro	Gly	Asp	Leu
				35					40					45
Arg	Asp	Leu	Gln	Val	Glu	Leu	Val	Lys	Thr	Ser	Val	Ala	Ala	Glu
				50					55					60
Glu	Phe	Ser	Ile	Leu	Met	Asn	Ile	Ser	Trp	Ile	Leu	Arg	Ala	Asp
				65					70					75
Ala	Ser	Ile	Arg	Leu	Leu	Lys	Ala	Thr	Lys	Ile	Cys	Val	Ser	Gly
				80					85					90
Lys	Asn	Asn	Met	Asn	Ser	Tyr	Ser	Cys	Val	Arg	Cys	Asn	Tyr	Thr
				95					100					105
Glu	Ala	Phe	Gln	Ser	Gln	Thr	Arg	Pro	Ser	Gly	Gly	Lys	Trp	Thr
				110					115					120
Phe	Ser	Tyr	Val	Gly	Phe	Pro	Val	Glu	Leu	Ser	Thr	Leu	Tyr	Leu
				125					130					135
Ile	Ser	Ala	His	Asn	Ile	Pro	Asn	Ala	Asn	Met	Asn	Glu	Asp	Ser
				140					145					150
Pro	Ser	Leu	Ser	Val	Asn	Phe	Thr	Ser	Pro	Gly	Cys	Leu	Asn	His
				155					160					165
Val	Met	Lys	Tyr	Lys	Lys	Gln	Cys	Thr	Glu	Ala	Gly	Ser	Leu	Trp
				170					175					180
Asp	Pro	Asp	Ile	Thr	Ala	Cys	Lys	Lys	Asn	Glu	Lys	Met	Val	Glu
				185					190					195
Val	Asn	Phe	Thr	Thr	Asn	Pro	Leu	Gly	Asn	Arg	Tyr	Thr	Ile	Leu
				200					205					210
Ile	Gln	Arg	Asp	Thr	Thr	Leu	Gly	Phe	Ser	Arg	Val	Leu	Glu	Asn
				215					220					225
Lys	Leu	Met	Arg	Thr	Ser	Val	Ala	Ile	Pro	Val	Thr	Glu	Glu	Ser
				230					235					240
Glu	Gly	Ala	Val	Val	Gln	Leu	Thr	Pro	Tyr	Leu	His	Thr	Cys	Gly
				245					250					255
Asn	Asp	Cys	Ile	Arg	Arg	Glu	Gly	Thr	Val	Val	Leu	Cys	Ser	Glu
				260					265					270
Thr	Ser	Ala	Pro	Ile	Pro	Pro	Asp	Asp	Asn	Arg	Arg	Met	Leu	Gly
				275					280					285
Gly	Trp	Leu	Pro	Leu	Phe	Leu	Val	Leu	Leu	Val	Ala	Val	Trp	Val
				290					295					300
Leu	Ala	Ala	Gly	Ile	Tyr	Leu	Thr	Trp	Arg	Gln	Gly	Arg	Ser	Thr
				305					310					315
Lys	Thr	Ser	Phe	Pro	Ile	Ser	Thr	Met	Leu	Leu	Pro	Leu	Ile	Lys
				320					325					330
Val	Leu	Val	Val	Tyr	Pro	Ser	Glu	Ile	Cys	Phe	His	His	Thr	Val
				335					340					345
Cys	Arg	Phe	Thr	Asp	Phe	Leu	Gln	Asn	Tyr	Cys	Arg	Ser	Glu	Val
				350					355					360
Ile	Leu	Glu	Lys	Trp	Gln	Lys	Lys	Lys	Ile	Ala	Glu	Met	Gly	Pro
				365					370					375
Val	Gln	Trp	Leu	Thr	Thr	Gln	Lys	Gln	Ala	Ala	Asp	Lys	Val	Val
				380					385					390
Phe	Leu	Leu	Pro	Ser	Asp	Val	Pro	Thr	Leu	Cys	Asp	Ser	Ala	Cys
				395					400					405
Gly	His	Asn	Glu	Gly	Ser	Ala	Arg	Glu	Asn	Ser	Gln	Asp	Leu	Phe
				410					415					420

SEQ 8/12

Pro	Leu	Ala	Phe	Asn	Leu	Phe	Cys	Ser	Asp	Phe	Ser	Ser	Gln	Thr
				425					430					435
His	Leu	His	Lys	Tyr	Leu	Val	Val	Tyr	Leu	Gly	Gly	Ala	Asp	Leu
				440					445					450
Lys	Gly	Asp	Tyr	Asn	Ala	Leu	Ser	Val	Cys	Pro	Gln	Tyr	His	Leu
				455					460					465
Met	Lys	Asp	Ala	Thr	Ala	Phe	His	Thr	Glu	Leu	Leu	Lys	Ala	Thr
				470					475					480
Gln	Ser	Met	Ser	Val	Lys	Lys	Arg	Ser	Gln	Ala	Cys	His	Asp	Ser
				485					490					495
Cys	Ser	Pro	Leu											
			499											

<210> 8
 <211> 289
 <212> PRT
 <213> Unknown

<220>

<221> PEPTIDE
 <223> Mouse Evi27 protein

Met	Leu	Leu	Val	Leu	Leu	Ile	Leu	Ala	Ala	Ser	Cys	Arg	Ser	Ala
				5					10					15
Leu	Pro	Arg	Glu	Pro	Thr	Ile	Gln	Cys	Gly	Ser	Glu	Thr	Gly	Pro
				20					25					30
Ser	Pro	Glu	Trp	Met	Val	Gln	His	Thr	Leu	Thr	Pro	Gly	Asp	Leu
				35					40					45
Arg	Asp	Leu	Gln	Val	Glu	Leu	Val	Lys	Thr	Ser	Val	Ala	Ala	Glu
				50					55					60
Glu	Phe	Ser	Ile	Leu	Met	Asn	Ile	Ser	Trp	Ile	Leu	Arg	Ala	Asp
				65					70					75
Ala	Ser	Ile	Arg	Leu	Leu	Lys	Ala	Thr	Lys	Ile	Cys	Val	Ser	Gly
				80					85					90
Lys	Asn	Asn	Met	Asn	Ser	Tyr	Ser	Cys	Val	Arg	Cys	Asn	Tyr	Thr
				95					100					105
Glu	Ala	Phe	Gln	Ser	Gln	Thr	Arg	Pro	Ser	Gly	Gly	Lys	Trp	Thr
				110					115					120
Phe	Ser	Tyr	Val	Gly	Phe	Pro	Val	Glu	Leu	Ser	Thr	Leu	Tyr	Leu
				125					130					135
Ile	Ser	Ala	His	Asn	Ile	Pro	Asn	Ala	Asn	Met	Asn	Glu	Asp	Ser
				140					145					150
Pro	Ser	Leu	Ser	Val	Asn	Phe	Thr	Ser	Pro	Gly	Cys	Leu	Asn	His
				155					160					165
Val	Met	Lys	Tyr	Lys	Lys	Gln	Cys	Thr	Glu	Ala	Gly	Ser	Leu	Trp
				170					175					180
Asp	Pro	Asp	Ile	Thr	Ala	Cys	Lys	Lys	Asn	Glu	Lys	Met	Val	Glu
				185					190					195
Val	Asn	Phe	Thr	Thr	Asn	Pro	Leu	Gly	Asn	Arg	Tyr	Thr	Ile	Leu
				200					205					210
Ile	Gln	Arg	Asp	Thr	Thr	Leu	Gly	Phe	Ser	Arg	Val	Leu	Glu	Asn
				215					220					225
Lys	Leu	Met	Arg	Thr	Ser	Val	Ala	Ile	Pro	Val	Thr	Glu	Glu	Ser
				230					235					240

SEQ 9/12

Glu	Gly	Ala	Val	Val	Gln	Leu	Thr	Pro	Tyr	Leu	His	Thr	Cys	Gly
				245					250					255
Asn	Asp	Cys	Ile	Arg	Arg	Glu	Gly	Thr	Val	Val	Leu	Cys	Ser	Glu
				260					265					270
Thr	Ser	Ala	Pro	Ile	Pro	Pro	Asp	Asp	Asn	Arg	Arg	Met	Leu	Gly
				275					280					285
Gly	Trp	Leu	Pro											
			289											

<210> 9
 <211> 866
 <212> PRT
 <213> Unknown

<220>

<221> PEPTIDE
 <223> IL-17 receptor protein

<400> 9														
Met	Gly	Ala	Ala	Arg	Ser	Pro	Pro	Ser	Ala	Val	Pro	Gly	Pro	Leu
				5					10					15
Leu	Gly	Leu	Leu	Leu	Leu	Leu	Leu	Gly	Val	Leu	Ala	Pro	Gly	Gly
				20					25					30
Ala	Ser	Leu	Arg	Leu	Leu	Asp	His	Arg	Ala	Leu	Val	Cys	Ser	Gln
				35					40					45
Pro	Gly	Leu	Asn	Cys	Thr	Val	Lys	Asn	Ser	Thr	Cys	Leu	Asp	Asp
				50					55					60
Ser	Trp	Ile	His	Pro	Arg	Asn	Leu	Thr	Pro	Ser	Ser	Pro	Lys	Asp
				65					70					75
Leu	Gln	Ile	Gln	Leu	His	Phe	Ala	His	Thr	Gln	Gln	Gly	Asp	Leu
				80					85					90
Phe	Pro	Val	Ala	His	Ile	Glu	Trp	Thr	Leu	Gln	Thr	Asp	Ala	Ser
				95					100					105
Ile	Leu	Tyr	Leu	Glu	Gly	Ala	Glu	Leu	Ser	Val	Leu	Gln	Leu	Asn
				110					115					120
Thr	Asn	Glu	Arg	Leu	Cys	Val	Arg	Phe	Glu	Phe	Leu	Ser	Lys	Leu
				125					130					135
Arg	His	His	His	Arg	Arg	Trp	Arg	Phe	Thr	Phe	Ser	His	Phe	Val
				140					145					150
Val	Asp	Pro	Asp	Gln	Glu	Tyr	Glu	Val	Thr	Val	His	His	Leu	Pro
				155					160					165
Lys	Pro	Ile	Pro	Asp	Gly	Asp	Pro	Asn	His	Gln	Ser	Lys	Asn	Phe
				170					175					180
Leu	Val	Pro	Asp	Cys	Glu	His	Ala	Arg	Met	Lys	Val	Thr	Thr	Pro
				185					190					195
Cys	Met	Ser	Ser	Gly	Ser	Leu	Trp	Asp	Pro	Asn	Ile	Thr	Val	Glu
				200					205					210
Thr	Leu	Glu	Ala	His	Gln	Leu	Arg	Val	Ser	Phe	Thr	Leu	Trp	Asn
				215					220					225
Glu	Ser	Thr	His	Tyr	Gln	Ile	Leu	Leu	Thr	Ser	Phe	Pro	His	Met
				230					235					240
Glu	Asn	His	Ser	Cys	Phe	Glu	His	Met	His	His	Ile	Pro	Ala	Pro
				245					250					255
Arg	Pro	Glu	Glu	Phe	His	Gln	Arg	Ser	Asn	Val	Thr	Leu	Thr	Leu
				260					265					270

SEQ 10/12

Arg	Asn	Leu	Lys	Gly	Cys	Cys	Arg	His	Gln	Val	Gln	Ile	Gln	Pro
				275					280					285
Phe	Phe	Ser	Ser	Cys	Leu	Asn	Asp	Cys	Leu	Arg	His	Ser	Ala	Thr
				290					295					300
Val	Ser	Cys	Pro	Glu	Met	Pro	Asp	Thr	Pro	Glu	Pro	Ile	Pro	Asp
				305					310					315
Tyr	Met	Pro	Leu	Trp	Val	Tyr	Trp	Phe	Ile	Thr	Gly	Ile	Ser	Ile
				320					325					330
Leu	Leu	Val	Gly	Ser	Val	Ile	Leu	Leu	Ile	Val	Cys	Met	Thr	Trp
				335					340					345
Arg	Leu	Ala	Gly	Pro	Gly	Ser	Glu	Lys	Tyr	Ser	Asp	Asp	Thr	Lys
				350					355					360
Tyr	Thr	Asp	Gly	Leu	Pro	Ala	Ala	Asp	Leu	Ile	Pro	Pro	Pro	Leu
				365					370					375
Lys	Pro	Arg	Lys	Val	Trp	Ile	Ile	Tyr	Ser	Ala	Asp	His	Pro	Leu
				380					385					390
Tyr	Val	Asp	Val	Val	Leu	Lys	Phe	Ala	Gln	Phe	Leu	Leu	Thr	Ala
				395					400					405
Cys	Gly	Thr	Glu	Val	Ala	Leu	Asp	Leu	Leu	Glu	Glu	Gln	Ala	Ile
				410					415					420
Ser	Glu	Ala	Gly	Val	Met	Thr	Trp	Val	Gly	Arg	Gln	Lys	Gln	Glu
				425					430					435
Met	Val	Glu	Ser	Asn	Ser	Lys	Ile	Ile	Val	Leu	Cys	Ser	Arg	Gly
				440					445					450
Thr	Arg	Ala	Lys	Trp	Gln	Ala	Leu	Leu	Gly	Arg	Gly	Ala	Pro	Val
				455					460					465
Arg	Leu	Arg	Cys	Asp	His	Gly	Lys	Pro	Val	Gly	Asp	Leu	Phe	Thr
				470					475					480
Ala	Ala	Met	Asn	Met	Ile	Leu	Pro	Asp	Phe	Lys	Arg	Pro	Ala	Cys
				485					490					495
Phe	Gly	Thr	Tyr	Val	Val	Cys	Tyr	Phe	Ser	Glu	Val	Ser	Cys	Asp
				500					505					510
Gly	Asp	Val	Pro	Asp	Leu	Phe	Gly	Ala	Ala	Pro	Arg	Tyr	Pro	Leu
				515					520					525
Met	Asp	Arg	Phe	Glu	Glu	Val	Tyr	Phe	Arg	Ile	Gln	Asp	Leu	Glu
				530					535					540
Met	Phe	Gln	Pro	Gly	Arg	Met	His	Arg	Val	Gly	Glu	Leu	Ser	Gly
				545					550					555
Asp	Asn	Tyr	Leu	Arg	Ser	Pro	Gly	Gly	Arg	Gln	Leu	Arg	Ala	Ala
				560					565					570
Leu	Asp	Arg	Phe	Arg	Asp	Trp	Gln	Val	Arg	Cys	Pro	Asp	Trp	Phe
				575					580					585
Glu	Cys	Glu	Asn	Leu	Tyr	Ser	Ala	Asp	Asp	Gln	Asp	Ala	Pro	Ser
				590					595					600
Leu	Asp	Glu	Glu	Val	Phe	Glu	Glu	Pro	Leu	Leu	Pro	Pro	Gly	Thr
				605					610					615
Gly	Ile	Val	Lys	Arg	Ala	Pro	Leu	Val	Arg	Glu	Pro	Gly	Ser	Gln
				620					625					630
Ala	Cys	Leu	Ala	Ile	Asp	Pro	Leu	Val	Gly	Glu	Glu	Gly	Gly	Ala
				635					640					645
Ala	Val	Ala	Lys	Leu	Glu	Pro	His	Leu	Gln	Pro	Arg	Gly	Gln	Pro
				650					655					660
Ala	Pro	Gln	Pro	Leu	His	Thr	Leu	Val	Leu	Ala	Ala	Glu	Glu	Gly
				665					670					675
Ala	Leu	Val	Ala	Ala	Val	Glu	Pro	Gly	Pro	Leu	Ala	Asp	Gly	Ala
				680					685					690

SEQ 11/12

Ala Val Arg Leu	Ala Leu Ala Gly Glu	Gly Glu Ala Cys Pro Leu	695	700	705
Leu Gly Ser Pro	Gly Ala Gly Arg Asn	Ser Val Leu Phe Leu Pro	710	715	720
Val Asp Pro Glu	Asp Ser Pro Leu Gly	Ser Ser Thr Pro Met Ala	725	730	735
Ser Pro Asp Leu	Leu Pro Glu Asp Val	Arg Glu His Leu Glu Gly	740	745	750
Leu Met Leu Ser	Leu Phe Glu Gln Ser	Leu Ser Cys Gln Ala Gln	755	760	765
Gly Gly Cys Ser	Arg Pro Ala Met Val	Leu Thr Asp Pro His Thr	770	775	780
Pro Tyr Glu Glu	Glu Gln Arg Gln Ser	Val Gln Ser Asp Gln Gly	785	790	795
Tyr Ile Ser Arg	Ser Ser Pro Gln Pro	Pro Glu Gly Leu Thr Glu	800	805	810
Met Glu Glu Glu	Glu Glu Glu Glu Gln	Asp Pro Gly Lys Pro Ala	815	820	825
Leu Pro Leu Ser	Pro Glu Asp Leu Glu	Ser Leu Pro Ser Leu Gln	830	835	840
Arg Gln Leu Leu	Phe Arg Gln Leu Gln	Lys Asn Ser Gly Trp Asp	845	850	855
Thr Met Gly Ser	Glu Ser Glu Gly Pro	Ser Ala	860	865	

PATENT COOPERATION TREATY

PCT

DECLARATION OF NON-ESTABLISHMENT OF INTERNATIONAL SEARCH REPORT

(PCT Article 17(2)(a), Rules 13ter and 39)

Applicant's or agent's file reference D6138PCT	IMPORTANT DECLARATION	Date of mailing (day/month/year) 06 JUN 2001
International application No. PCT/US01/03518	International filing date (day/month/year) 02 FEBRUARY 2001	(Earliest) Priority Date (day/month/year) 04 FEBRUARY 2000
International Patent Classification (IPC) or both national classification and IPC Please See Continuation Sheet.		
Applicant THE BOARD OF TRUSTEES OF THE UNIVERSITY OF ARKANSAS		

This International Searching Authority hereby declares, according to Article 17(2)(a), that **no international search report will be established on the international application for the reasons indicated below.**

1. ☒ The subject matter of the international application relates to:
 - a. ☐ scientific theories.
 - b. ☐ mathematical theories.
 - c. ☐ plant varieties.
 - d. ☒ animal varieties.
 - e. ☐ essentially biological processes for the production of plants and animals, other than microbiological processes and the products of such processes.
 - f. ☐ schemes, rules or methods of doing business.
 - g. ☐ schemes, rules or methods of performing purely mental acts.
 - h. ☐ schemes, rules or methods of playing games.
 - i. ☐ methods for treatment of the human body by surgery or therapy.
 - j. ☐ methods for treatment of the animal body by surgery or therapy.
 - k. ☐ diagnostic methods practiced on the human or animal body.
 - l. ☐ mere presentations of information.
 - m. ☐ computer programs for which this International Searching Authority is not equipped to search prior art.
2. ☐ The failure of the following parts of the international application to comply with prescribed requirements prevents a meaningful search from being carried out:

☐ the description
 ☐ the claims
 ☐ the drawings
3. ☒ The failure of the nucleotide and/or amino acid sequence listing to comply with the standard provided for in Annex C of the Administrative Instructions prevents a meaningful search from being carried out.

☐ the written form has not been furnished or does not comply with the standard.
 ☒ the computer readable form has not been furnished or does not comply with the standard.
4. Further comments:
Please See Continuation Sheet.

Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer <div style="text-align: center;"> TERRY J. DEY PARALEGAL SPECIALIST TECHNOLOGY CENTER 1600 </div> DONG JIANG Telephone No. (703) -308-0196
---	--

Form PCT/ISA/203 (July 1998)★

**DECLARATION OF NON-ESTABLISHMENT OF
INTERNATIONAL SEARCH REPORT**

International application No.
PCT/US01/03518

The International Patent Classification (IPC) or National Classification and IPC are as listed below:

IPC(7): C12N 15/00; C07K 1/00
US Cl.: 435/69.1, 7.1; 530/350, 387.9,

4. Further Comments (Continued):

Although the claims contain other descriptions besides SEQ ID NOs, such as a name for the claimed protein (Evi27), the molecular weight ("about 24, 33, 56, 47, 75, 127, and 150 kD", claim 9), and certain biological activity (stimulating the secretion of IL-1, IL-18, and TNF- α by hematopoietic cells, claims 17-19), they are not meaningful for conducting a specific search of the prior art, and obtaining pertinent results for the following reasons:

1. Claims 17, and 20-21 describe the polypeptide of interest by an arbitrary name, Evi27, which does not confer the distinct structural identity of the protein. Additionally, the same protein isolated by others, may be given different names, and other proteins with distinct structural and functional properties may be described by the same name. Therefore, a name alone is not meaningful.
2. Claim 9 describes said protein by molecular weight, which, by itself, is not sufficient to distinguish proteins as one or the other [credible] because distinct proteins can have the same molecular weight, and different molecular weight can be a result of multiple reasons, such as post-translational modification, or protease effect.
3. The functional activity of the protein asserted in claims 17-19 is not specific. There are known other factors that can stimulate hematopoietic cells to secrete IL-1, IL-18, and TNF- α .

THIS PAGE BLANK (USPTO)